arXiv:math/0010250v1 [math.QA] 26 Oct 2000

On FRT– Clifford Algebras

I. Heckenberger A. Schüler*

Universität Leipzig, Mathematisches Institut,
Augustusplatz 10–11, 04109 Leipzig, Germany
e-mail:heckenbe,schueler@mathematik.uni-leipzig.de

Abstract

We study the q-Clifford algebras $\mathrm{Cl}_{q^2}(N, c^2)$, called FRT– Clifford algebras, introduced by Faddeev, Reshetikhin and Takhtajan. It is shown that $\mathrm{Cl}_{q^2}(N, c^2)$ acts on the q-exterior algebra $\Lambda(\mathrm{O}_{q^2}^N)$. Moreover, explicit formulas for the embedding of $U_{q^2}(\mathfrak{so}_N)$ into $\mathrm{Cl}_{q^2}(N, c^2)$ and its relation to the vector and spin representations of $U_{q^2}(\mathfrak{so}_N)$ are given and proved.

MSC 2000: 81R50, 15A66

Key Words: q-Clifford algebra, Drinfeld-Jimbo algebra, spin representation

1 Introduction

The theory of spinors gives one of the most powerful structures in differential geometry and theoretical physics. Its origin can be traced back to works of Cartan, Dirac, Brauer and Weyl. On the other hand, quantum groups and quantum spaces appeared in the eighties as quantized algebras of functions on complex manifolds. Because of their close relation to Lie groups and homogeneous spaces one can ask whether one can develop a spin geometry of quantum spaces. In the last years there were made several attempts in this direction. Quantum Clifford algebras are introduced and studied for instance in [11], [4], [10], [1], [9].

In [19] Woronowicz presented a theory of covariant differential calculus over Hopf algebras. Using this or related theories quantum Clifford algebras and spinors are defined and investigated by many authors, see [8], [6], [7], [2], [14]. The approach of Woronowicz causes two important facts which are absent in the classical situation. First, in general there exists no canonical differential calculus on a given quantum space. Given a differential calculus, the corresponding external algebra of differential forms depends extremely on the algebra of functions and on the first order calculus. Secondly, because of the noncommutative structure defining properties of a metric tensor are much more restrictive than in the commutative situation. Hence the varieties of Clifford algebras and spinors in the classical and quantum setting are very different.

---

*both authors are supported by the Deutsche Forschungsgemeinschaft

Exterior algebras $\Lambda(\mathrm{O}_{q^2}^N)$ of the quantum Euclidean space $\mathrm{O}_{q^2}(N)$ are introduced in [8], [17] and [12]. They naturally appeared in [5] and [18] where the higher order calculi of the coordinate algebra of the quantum Euclidean space and the quantum Euclidean sphere, respectively, were examined. The corresponding quantum Clifford algebras, called FRT– Clifford algebras, are defined in [8] and analyzed in [6]. The aim of this paper is to prove further results on FRT– Clifford algebras and to present elementary methods for the dealing with these objects. We construct spin representations of $U_{q^2}(\mathfrak{so}_N)$ using FRT– Clifford algebras. Related quantum spin bundles on quantum spaces will be examined in a forthcoming paper.

In Section 2 the definition of the FRT– Clifford algebra $\mathrm{Cl}_{q^2}(N,c^2)$ is recalled. In Theorem 1 we prove that the algebra $\mathrm{Cl}_{q^2}(N,c^2)$ has a representation over the quantum exterior algebra $\Lambda(\mathrm{O}_{q^2}^N)$. Elementary proofs of the semisimplicity of the algebras $\mathrm{Cl}_{q^2}(N,c^2)$ are given. Canonical minimal left- and right-ideals of $\mathrm{Cl}_{q^2}(N,c^2)$ are introduced. In Section 3 we formulate and give the proof of two theorems. In Theorem 8 an algebra map π from $U_{q^2}(\mathfrak{so}_N)$ to $\mathrm{Cl}_{q^2}(N,c^2)$ is given. In Theorem 9 invariance of the vector space of generators of $\mathrm{Cl}_{q^2}(N,c^2)$ under the adjoint action of $\pi(U_{q^2}(\mathfrak{so}_N))$ is proved. Further, the spin representations of $U_{q^2}(\mathfrak{so}_N)$ are realized on the canonical left ideal of $\mathrm{Cl}_{q^2}(N,c^2)$.

For the set of nonzero real and complex numbers we take the symbols $\mathbb{R}^\times$ and $\mathbb{C}^\times$, respectively. Throughout we use Einsteins convention to sum over repeated indices. We write $\delta^i_j = \delta_{ij}$ for Kronecker's symbol. For integer numbers j, $1 \le j \le N$, the symbol j' means the number $N+1-j$. For $m, p \in \mathbb{N}_0$ we set

$$[\![m]\!] = \frac{1-q^{-4m}}{1-q^{-4}}, \quad [\![m]\!]! = \prod_{k=1}^{m} [\![k]\!], \quad [\![0]\!]! = 1, \quad \begin{bmatrix}\!\begin{bmatrix} m \\ p \end{bmatrix}\!\end{bmatrix} = \frac{[\![m]\!]!}{[\![p]\!]![\![m-p]\!]!}. \tag{1}$$

We use the definition of the Birman–Wenzl–Murakami algebra $\mathrm{BWM}(q^{N-1},q)_k$ with $2k-2$ generators g_i, e_i, $i=1,2,\dots,k-1$, given in [3], in the notation of [15]. The vector space of intertwiners of corepresentations v,v' of the coordinate Hopf algebra $\mathcal{O}(\mathrm{O}_{q^2}(N))$ of the quantum group $\mathrm{O}_{q^2}(N)$ is denoted by $\mathrm{Mor}(v,v')$. Let u denote the fundamental (vector) corepresentation of $\mathcal{O}(\mathrm{O}_{q^2}(N))$. For the generators of $\mathrm{Mor}(u\otimes u,u\otimes u)$, $\mathrm{Mor}(1,u\otimes u)$ and $\mathrm{Mor}(u\otimes u,1)$ we use the following symbols in the graphical calculus (see [5]):

$\hat{R}=$  $\hat{R}^{-1}=$  $K=$  $C^{ij}=$  $C_{ij}=$

2 The FRT– Clifford algebras

Let q be a nonzero complex number, $q^m\neq 1$ for all $m\in\mathbb{N}$. We set $[2]=q+q^{-1}$ and $\hat{q}=q-q^{-1}$. Let $N\in\mathbb{N}$, $N\geq 3$, and $n\in\mathbb{N}$ such that $N=2n+\epsilon$, $\epsilon\in\{0,1\}$. In the sequel we frequently use the following standard notation for $\hat{R}$-matrices corresponding to orthogonal quantum groups $\mathrm{O}_{q^2}(N)$, cf. [8, Subsect. 1.4].

$$\begin{gathered}\hat{R}^{ij}_{kl}=q^{2\delta^i_j-2\delta^i_{j'}}\delta^i_l\delta^j_k+(q^2-q^{-2})\delta_{i<l}(\delta^i_k\delta^j_l-K^{ij}_{kl}),\\ C^{ij}=C_{ij}=q^{-2\rho_i}\delta^i_{j'},\quad K^{ij}_{kl}=C^{ij}C_{kl},\qquad \rho_i=\delta_{i<i'}(N/2-i)-\delta_{i'<i}(N/2-i'),\\ P_+=\frac{1}{q^2+q^{-2}}\left(q^{-2}\,\mathrm{id}+\hat{R}-\frac{[2]\hat{q}}{q^{2N}-1}K\right),\\ P_-=\frac{1}{q^2+q^{-2}}\left(q^{2}\,\mathrm{id}-\hat{R}-\frac{[2]\hat{q}}{q^{2N-4}+1}K\right),\end{gathered}\tag{2}$$

where $\delta_{i<l}=1$ if $i<l$ and $\delta_{i<l}=0$ otherwise.

Let $c^2\in\mathbb{R}^\times$. By Definition 13 in [8] the quantum Clifford algebra $\mathrm{Cl}_{q^2}(N,c^2)$ is the complex unital algebra generated by the elements γ_i, $i=1,\dots,N$, and relations

$$P_{+}{}^{kl}_{ij}\gamma_k\gamma_l=0,\quad i,j=1,2,\dots,N,\qquad\qquad C^{ij}\gamma_i\gamma_j=c^2\frac{q^{2N}-1}{q^2-1}.\tag{3}$$

The factor in the last relation is choosen in such a way that formulas in the sequel become more simple. The explicit form of the relations of $\mathrm{Cl}_{q^2}(N,c^2)$ can be computed similarly to the proof of Proposition 9.14 in [16]. We obtain

$$
\begin{gathered}
\gamma_i\gamma_i = 0, \quad i \neq i', \qquad \gamma_j\gamma_i = -q^2\gamma_i\gamma_j, \quad j > i, j \neq i', \\
\gamma_{i'}\gamma_i = -\gamma_i\gamma_{i'} + (q^2 - q^{-2})\sum_{j=1}^{i-1} q^{2j-2i+2}\gamma_j\gamma_{j'} + c^2 q^{N-2i+1}[2], \quad i < i', \\
\epsilon\gamma_{n+1}\gamma_{n+1} = \epsilon\left(\hat{q}\sum_{j=1}^{n} q^{2j-2n}\gamma_j\gamma_{j'} + c^2\right).
\end{gathered} \tag{4}
$$

It is easy to see that the mapping $\gamma_i \mapsto c\gamma_i$ defines an algebra isomorphism $\mathrm{Cl}_{q^2}(N, c^2) \to \mathrm{Cl}_{q^2}(N, 1)$. In contrast, the choice $c = 0$ leads to the q-exterior algebra $\Lambda(\mathrm{O}_{q^2}^N)$ which has been studied for instance in [8], [17] or [12, Sect. 5].

The algebra $\mathrm{Cl}_{q^2}(N, c^2)$ obeys a $\mathbb{Z}_2$-grading ∂_0 given by $\partial_0(\gamma_i) = 1$ for all $i = 1, \dots, N$. The components of even and odd degree are denoted by $\mathrm{Cl}_{q^2}^{+}(N, c^2)$ and $\mathrm{Cl}_{q^2}^{-}(N, c^2)$, respectively. There exists also a $\mathbb{Z}^n$-grading $\partial = (\partial_1, \partial_2, \dots, \partial_n)$ of $\mathrm{Cl}_{q^2}(N, c^2)$ which is determined by $\partial_i(\gamma_j) = \delta_j^i - \delta_{j'}^i$ for $i = 1, \dots, n$ and $j = 1, \dots, N$. For $N = 2n+1$ the mapping $\partial_{n+1} : \gamma_j \mapsto \delta_{n+1}^j$, $j = 1, 2, \dots, N$, defines another $\mathbb{Z}_2$-grading of $\mathrm{Cl}_{q^2}(2n+1, c^2)$.

Let α_i, $i = 1, \dots, N$, be nonzero complex numbers such that $\alpha_i\alpha_{i'} = 1$. From (4) it follows that the mapping $\gamma_i \mapsto \alpha_i\gamma_i$ defines an algebra isomorphism of $\mathrm{Cl}_{q^2}(N, c^2)$. Moreover, there exists an algebra antiautomorphism τ of $\mathrm{Cl}_{q^2}(N, c^2)$ given by $\tau(\gamma_i) = \gamma_{i'}$. This antiautomorphism as well as its compose with any of the above automorphisms has order two.

Let V denote the complex vector space with basis $\{\gamma_i \,|\, i = 1, \dots, N\}$ and let $V^{\otimes k}$, $k \geq 1$, be the k-fold tensor product $V \otimes V \otimes \cdots \otimes V$. We write $V^{\otimes}$ for $\bigoplus_{k=0}^{\infty} V^{\otimes k}$, where $V^{\otimes 0} = \mathbb{C}1$. The mapping $\varphi : \mathrm{BWM}(q^{2N-2}, q^2)_{k+1} \to \mathrm{End}(V^{\otimes k+1})$, given by $\varphi(g_i) = \hat{R}_{i,i+1}$, $\varphi(e_i) = K_{i,i+1}$, $i = 1, \dots, k$, is a representation of $\mathrm{BWM}(q^{2N-2}, q^2)_{k+1}$. If no confusion can arise we will identify elements of the algebra $\mathrm{BWM}(q^{2N-2}, q^2)_{k+1}$ with their images under the representation φ.

In [15], formulas (19) and (20), the elements ${d'}_{k+1,i}^{-}$ and $b_{1,k}^{-}$, $k = 0, 1, \dots, N-1$, $i = 1, \dots, k$, of the Birman–Wenzl–Murakami algebra $\mathrm{BWM}(q^{2N-2}, q^2)_{k+1}$ were introduced.

$$
\begin{gathered}
{d'}_{k+1,i}^{-} = K_{12}K_{23}\cdots K_{k-i+1,k-i+2}\sum_{j=0}^{i-1}(-q^{-2})^j\prod_{l=1}^{j}\hat{R}_{k-i+1+l,k-i+2+l}, \\
b_{1,k}^{-} = \sum_{i=0}^{k}(-q^{-2})^i\hat{R}_{12}\hat{R}_{23}\cdots\hat{R}_{i,i+1} - \frac{[2]\hat{q}}{1+q^{2N-4k}}\sum_{i=1}^{k} q^{4i-4k-2}{d'}_{k+1,i}^{-}.
\end{gathered} \tag{5}
$$

Using (5) the unique antisymmetrizer A_{k+1} of $\mathrm{BWM}(q^{2N-2},q^2)_{k+1}$ is constructed. Recall that $A_1=\mathrm{id}$, $A_2=P_-$ and $A_{i+1}=(\mathrm{id}\otimes A_i)b^-_{1,i}/\llbracket i+1\rrbracket$ for $i>2$. Further, from $A_{m+1}=A_{m+1}(\mathrm{id}\otimes A_m)$, $\hat{R}_{i,i+1}A_{m+1}=-q^{-2}A_{m+1}$, $K_{i,i+1}A_{m+1}=0$ $(i=1,2,\dots,m)$ and (5) we conclude that

$$\begin{aligned} A_{m+1}&=\frac{1}{\llbracket m+1\rrbracket}(\mathrm{id}\otimes A_m)b^-_{1,m}=\frac{1}{\llbracket m+1\rrbracket}(\mathrm{id}\otimes A_m)b^-_{1,m}(\mathrm{id}\otimes A_m)\\ &=\frac{1}{\llbracket m+1\rrbracket}(\mathrm{id}\otimes A_m)\left(\mathrm{id}-q^{-2}\llbracket m\rrbracket\hat{R}_{12}-\frac{1-q^{-4m}}{1+q^{2N-4m}}K_{12}\right)(\mathrm{id}\otimes A_m). \end{aligned}\tag{6}$$

We use the symbol $V^{\wedge k}$ for the vector space $V^{\otimes k}/\ker A_k$. Then $\Lambda(\mathrm{O}^N_{q^2})=\bigoplus_{k=0}^N V^{\wedge k}$ as vector spaces and the multiplication $\wedge$ corresponds to the tensor product in $V^{\otimes}$.

Let $g:V\otimes V\to\mathbb{C}$ denote the linear mapping given by the formula $g(\gamma_i\otimes\gamma_j)=c^2q^{2N-3}[2]C_{ij}/(q^{2N-4}+1)$. From $C^{ij}C_{ij}=q^{-2N+2}(q^{2N}-1)(q^{2N-4}+1)/(q^2-q^{-2})$ we then get $a_{ij}\gamma_i\gamma_j=g(a_{ij}\gamma_i\otimes\gamma_j)$ in $\mathrm{Cl}_{q^2}(N,c^2)$ whenever $a_{ij}\gamma_i\wedge\gamma_j=0$ in $\Lambda(\mathrm{O}^N_{q^2})$. Inspired by the methods presented in [13] and [2], let us define contraction mappings $\langle\!\langle\cdot,\cdot\rangle\!\rangle:V^{\otimes k}\otimes V^{\otimes l}\to V^{\otimes\max\{0,l-k\}}$ by $\langle\!\langle\rho_k,\rho'_l\rangle\!\rangle=0$ for $k>l$ and

$$\langle\!\langle\rho,\rho'_l\rangle\!\rangle=g_{12}(\rho\otimes b^-_{1,l-1}(\rho'_l)),\qquad\qquad \langle\!\langle\rho\otimes\rho_k,\rho'_l\rangle\!\rangle=\langle\!\langle\rho,\langle\!\langle\rho_k,\rho'_l\rangle\!\rangle\rangle\!\rangle\tag{7}$$

for $\rho\in V$, $\rho_i,\rho'_i\in V^{\otimes i}$. Now let $l\geq k$, $\rho'_l\in V^{\otimes l}$, $\rho_k:=a_J\gamma_{j_1}\gamma_{j_2}\cdots\gamma_{j_k}\in V^{\otimes k}$, $a_J\in\mathbb{C}$ for $J=(j_1,\dots,j_k)$, $1\leq j_m\leq N$ for all m. We show that $A_{l-k}\langle\!\langle\rho_k,\rho'_l\rangle\!\rangle=0$ provided $A_l(\rho'_l)=0$ or $A_k(\rho_k)=0$. Indeed, by the definition of the contraction mapping we get

$$\begin{aligned} A_{l-k}\langle\!\langle\rho_k,\rho'_l\rangle\!\rangle&=a_JA_{l-k}\langle\!\langle\gamma_{j_1},\langle\!\langle\gamma_{j_2},\cdots\langle\!\langle\gamma_{j_k},\rho'_l\rangle\!\rangle\rangle\!\rangle\rangle\!\rangle\\ &=a_Jg_{12}(\gamma_{j_1}\otimes\llbracket l-k+1\rrbracket A_{l-k+1}(\langle\!\langle\gamma_{j_2},\cdots\langle\!\langle\gamma_{j_k},\rho'_l\rangle\!\rangle\rangle\!\rangle))=\cdots\\ &=\frac{\llbracket l\rrbracket!}{\llbracket l-k\rrbracket!}a_Jg_{12}(\gamma_{j_1}\otimes g_{12}(\gamma_{j_2}\otimes\cdots g_{12}(\gamma_{j_k}\otimes A_l(\rho'_l)))). \end{aligned}$$

Hence $\langle\!\langle\cdot,\cdot\rangle\!\rangle$ maps $V^{\otimes k}\otimes V^{\wedge l}$, $l\geq k$, to $V^{\wedge l-k}$. On the other hand, the last line of the above equation can also be written in the form

$$A_{l-k}\langle\!\langle\rho_k,\rho'_l\rangle\!\rangle=\frac{\llbracket l\rrbracket!}{\llbracket l-k\rrbracket!}g_{12}g_{23}\cdots g_{k,k+1}(\rho_k\otimes A_l(\rho'_l)).$$

Suppose now that $A_k\rho_k=0$. Then ρ_k is a linear combination of elements $\xi_0:=\xi'_i\otimes\xi\otimes\xi''_{k-i-2}$, $\xi'_i\in V^{\otimes i}$, $\xi''_{k-i-2}\in V^{\otimes k-i-2}$, $\xi\in V\otimes V$, $A_2(\xi)=0$. Without loss of generality we can assume that $\hat{R}\xi=r\xi$ with $r=q^2$ or $r=q^{-2N+2}$. This means that ξ is an element of the image of the projector P_+ or P_0, respectively. For such elements we easily obtain

$$\begin{aligned}
A_{l-k}\langle\!\langle \xi_0, \rho_l' \rangle\!\rangle &= A_{l-k}\langle\!\langle r^{-1}\hat{R}_{i+1,i+2}\xi_0, \rho_l' \rangle\!\rangle \\
&= \frac{r^{-1}[\![l]\!]!}{[\![l-k]\!]!} g_{12}g_{23}\cdots g_{k,k+1}(\hat{R}_{i+1,i+2}\xi_0 \otimes A_l(\rho_l')) \\
&= \frac{r^{-1}[\![l]\!]!}{[\![l-k]\!]!} g_{12}g_{23}\cdots g_{k,k+1}(\xi_0 \otimes \hat{R}_{k-i-1,k-i}A_l(\rho_l')) \\
&= \frac{-q^{-2}r^{-1}[\![l]\!]!}{[\![l-k]\!]!} g_{12}g_{23}\cdots g_{k,k+1}(\xi_0 \otimes A_l(\rho_l')) = -q^{-2}r^{-1}A_{l-k}\langle\!\langle \xi_0, \rho_l' \rangle\!\rangle.
\end{aligned}$$

Since q is not a root of unity, we obtain $A_{l-k}\langle\!\langle \xi_0, \rho_l' \rangle\!\rangle = 0$ and hence $A_{l-k}\langle\!\langle \rho_k, \rho_l' \rangle\!\rangle = 0$. This means that $\langle\!\langle \cdot, \cdot \rangle\!\rangle$ is a well-defined mapping from $V^{\wedge k} \otimes V^{\wedge l}$, $k \le l$, to $V^{\wedge l-k}$.

Theorem 1. *There exists a representation* $\triangleright$ *of* $\mathrm{Cl}_{q^2}(N, c^2)$ *on* $\Lambda(\mathrm{O}_{q^2}^N)$ *such that*

$$\gamma_i \triangleright \rho_m = \gamma_i{\wedge}\rho_m + \frac{1+q^{2N-4}}{1+q^{2N-4m}}\langle\!\langle \gamma_i, \rho_m \rangle\!\rangle \tag{8}$$

for $\rho_m \in V^{\wedge m}$, $m \in \{0, 1, \ldots, N\}$, $i = 1, 2, \ldots, N$.

Proof. Since $\langle\!\langle \cdot, \cdot \rangle\!\rangle$ is a well-defined mapping from $V \otimes V^{\wedge m}$ to $V^{\wedge m-1}$, we only have to show that the definition (8) is compatible with the relations (3) of $\mathrm{Cl}_{q^2}(N, c^2)$. Suppose that $m = 0$. Then

$$\gamma_i\gamma_j \triangleright 1 = \gamma_i \triangleright \gamma_j = \gamma_i{\wedge}\gamma_j + \langle\!\langle \gamma_i, \gamma_j \rangle\!\rangle$$

for $i, j = 1, \ldots, N$. Since $g(P_{+}{}^{kl}_{ij}\gamma_k \otimes \gamma_l) = 0$ we obtain $P_{+}{}^{kl}_{ij}\gamma_k\gamma_l \triangleright 1 = 0$. From $C^{ij}C_{ij} = q^{-2N+2}(q^{2N}-1)(q^{2N-4}+1)/(q^2-q^{-2})$ we conclude that

$$C^{ij}\gamma_i\gamma_j \triangleright 1 = C^{ij}g(\gamma_i \otimes \gamma_j) = C^{ij}\frac{c^2q^{2N-3}[2]}{q^{2N-4}+1}C_{ij} = \frac{c^2q^{-1}(q^{2N}-1)}{\hat{q}}$$

which proves the compatibility with the second equation of (3) for $m = 0$.

Now consider the case $m \ge 1$. Then we get

$$\begin{aligned}
a_{ij}\gamma_i\gamma_j \triangleright \rho_m &= a_{ij}\gamma_i \triangleright \left(\gamma_j{\wedge}\rho_m + \frac{1+q^{2N-4}}{1+q^{2N-4m}}\langle\!\langle \gamma_j, \rho_m \rangle\!\rangle \right) \\
&= a_{ij}\gamma_i{\wedge}\gamma_j{\wedge}\rho_m + \frac{(1+q^{2N-4})^2}{(1+q^{2N-4m})(1+q^{2N-4m+4})}\langle\!\langle a_{ij}\gamma_i \otimes \gamma_j, \rho_m \rangle\!\rangle \\
&\quad + \frac{1+q^{2N-4}}{1+q^{2N-4m}}a_{ij}\gamma_i{\wedge}\langle\!\langle \gamma_j, \rho_m \rangle\!\rangle + \frac{1+q^{2N-4}}{1+q^{2N-4m-4}}a_{ij}\langle\!\langle \gamma_i, \gamma_j \otimes \rho_m \rangle\!\rangle
\end{aligned}$$

for $\rho_m \in V^{\wedge m}$, $i, j = 1, \dots, N$, $a_{ij} \in \mathbb{C}$. If $a_{ij}\gamma_i \otimes \gamma_j \in \ker A_2$ then the first two summands of the last equation vanish and we get an element in $V^{\wedge m}$. More precisely, for $\rho_m = \gamma_{k_1}\wedge\gamma_{k_2}\wedge\cdots\gamma_{k_m}$ the expression $A_m(a_{ij}\gamma_i\gamma_j \triangleright \rho_m)$ takes the form

$$\frac{a_{ij}c^2q^{2N-3}[2]}{1+q^{2N-4}}\left(\frac{1+q^{2N-4}}{1+q^{2N-4m}}\right.$$



$$+\frac{1+q^{2N-4}}{1+q^{2N-4m-4}}$$



$$\left.\vphantom{\frac{1}{1}}\right)\gamma_{l_1}\wedge\gamma_{l_2}\wedge\cdots\wedge\gamma_{l_m}.$$

Using $A_m = A_m(\mathrm{id}\otimes A_{m-1})$ and $(\mathrm{id}\otimes A_{m-1})b^-_{1,m-1} = \llbracket m\rrbracket A_m$ in the first summand and $(\mathrm{id}\otimes A_m)b^-_{1,m} = \llbracket m+1\rrbracket A_{m+1}$ in the second one the coefficient in the above expression becomes

$$a_{ij}\left(\frac{c^2q^{2N-3}[2]\llbracket m\rrbracket}{1+q^{2N-4m}}\right.$$



$$+\frac{c^2q^{2N-3}[2]\llbracket m+1\rrbracket}{1+q^{2N-4m-4}}$$



$$\left.\vphantom{\frac{1}{1}}\right). \qquad (*)$$

Now in the second summand we replace A_{m+1} by the second line of (6). Then for $a_{ij} = C^{ij}$ the expression $(*)$ becomes

$$\left(\frac{c^2q^{2N-3}[2]\llbracket m\rrbracket}{q^{2N-4m}+1} + \frac{c^2q^{2N-3}[2]}{q^{2N-4m-4}+1}\left(\frac{q^{-2N+2}(q^{2N}-1)(q^{2N-4}+1)}{[2]\hat{q}} - \right.\right.$$
$$\left.\left. - q^{-2}\llbracket m\rrbracket q^{2N-2} - \frac{1-q^{-4m}}{1+q^{2N-4m}}\right)\right)A_m = \frac{c^2(q^{2N}-1)}{q^2-1}A_m$$

which proves the compatibility of $\triangleright$ with the last relation in (3). Setting $a_{ij} = P_{+}{}^{ij}_{rs}$, $r, s = 1, \dots, N$, $(*)$ results in

$$\left(\frac{c^2q^{2N-3}[2]\llbracket m\rrbracket}{q^{2N-4m}+1} + \frac{c^2q^{2N-3}[2]}{q^{2N-4m-4}+1}\left(0 - q^{-2}\llbracket m\rrbracket q^{-2} - \frac{1-q^{-4m}}{1+q^{2N-4m}}\right)\right)M_{+,m} = 0,$$

where

$$M_{+,m} =$$



.

■

Let $I_N = (i_1, i_2, \dots, i_N)$ and $J_n = (j_1, j_2, \dots, j_n)$, $i_k, j_k \in \{0,1\}$ for all k. We use the abbreviation $\gamma^{I_N} = \gamma_1^{i_1}\gamma_2^{i_2}\cdots\gamma_N^{i_N}$. Set $\ell(I_N) = \sum_{k=1}^N i_k$, $\ell(J_n) = \sum_{k=1}^n j_k$, and $L(J_n) = \sum_{k=1}^n k j_k$.

Proposition 2. *The set* $\{\gamma_1^{i_1}\gamma_2^{i_2}\cdots\gamma_N^{i_N} \,|\, i_k \in \{0,1\}, k = 1,\dots,N\}$ *forms a vector space basis of* $\mathrm{Cl}_{q^2}(N, c^2)$.

Proof. From the explicit form (4) of the generating relations of the algebra $\mathrm{Cl}_{q^2}(N,c^2)$ we conclude that the elements γ^{I_N} span the vector space $\mathrm{Cl}_{q^2}(N,c^2)$. To prove that they form a basis we use Theorem 1. Suppose that there is an element $\rho = \sum_{I_N} \lambda_{I_N}\gamma^{I_N} = 0$ in $\mathrm{Cl}_{q^2}(N,c^2)$. Then $\rho \triangleright 1 = 0$. Let $j_{\max} = \max\{\ell(I_N) \,|\, \lambda_{I_N} \neq 0\}$. Then we have

$$\rho \triangleright 1 = \sum_{J_N, \ell(J_N) = j_{\max}} \lambda_{J_N} \gamma_1^{j_1} \wedge \gamma_2^{j_2} \wedge \cdots \wedge \gamma_N^{j_N} + \rho',$$

where $\rho' \in \bigoplus_{k=0}^{j_{\max}-1} V^{\wedge k}$. Since $\rho \triangleright 1 = 0$, we conclude that $\lambda_{J_N} = 0$ for all J_N with $\ell(J_N) = j_{\max}$. This is a contradiction to the definition of $j_{\max}$. ■

Corollary 3. *The representation* $\triangleright$ *of* $\mathrm{Cl}_{q^2}(N,c^2)$ *on* $\Lambda(\mathrm{O}_{q^2}^N)$ *is faithful.*

Let $\widehat{\mathrm{Cl}}_{q^2}(N,c^2)$ denote the subalgebra of $\mathrm{Cl}_{q^2}(N,c^2)$ defined by

$$\widehat{\mathrm{Cl}}_{q^2}(N,c^2) = \{z \in \mathrm{Cl}_{q^2}(N,c^2) \,|\, \partial_k(z) = 0 \quad \text{for all } k = 1,2,\dots,n\}. \tag{9}$$

Proposition 4. *The algebra* $\widehat{\mathrm{Cl}}_{q^2}(N,c^2)$ *is commutative.*

Proof. By Proposition 2 the set

$$\{\gamma_1^{i_1}\gamma_2^{i_2}\cdots\gamma_n^{i_n}\gamma_{n+1}^{i_{n+1}}\gamma_{n'}^{i_n}\cdots\gamma_{N-1}^{i_2}\gamma_N^{i_1} \,|\, i_l \in \{0,1\}, l = 1,2,\dots,n+1,\ i_{n+1} \le \epsilon\} \tag{10}$$

forms a vector space basis of the algebra $\widehat{\mathrm{Cl}}_{q^2}(2n+\epsilon, c^2)$. From this fact and the first line in (4) we conclude that $\widehat{\mathrm{Cl}}_{q^2}(N,c^2)$ is generated as an algebra by the set of its elements

$$\begin{aligned} &\{\gamma_i\gamma_{i'} \,|\, i = 1,2,\dots,n\} && \text{for } N = 2n,\\ &\{\gamma_{n+1}, \gamma_i\gamma_{i'} \,|\, i = 1,2,\dots,n\} && \text{for } N = 2n+1. \end{aligned}$$

Now the commutativity of these generators follows immediately from the equations in the first line of (4). ■

The following proposition, as well as Proposition 6 was proved in [6] using representation theory of $U_{q^2}(\mathfrak{so}_N)$. Here we give an elementary proof.

Proposition 5. *The algebra* $\mathrm{Cl}_{q^2}(2n, c^2)$ *is simple.*

Proof. Let $\mathcal{I} \neq \{0\}$ be a two-sided ideal of $\mathrm{Cl}_{q^2}(2n, c^2)$. We prove that $\mathcal{I} = \mathrm{Cl}_{q^2}(2n, c^2)$. First it will be shown by induction that

($*$) for all $j \in \{1, 2, \dots, n\}$ there exists $\rho_j = \sum_{I_N} \lambda_{I_N} \gamma^{I_N} \in \mathcal{I}$ such that $\lambda_{I_N} \neq 0$ implies $i_k = 1$ for all $k < j$.

For $j = 1$ this only means that there exists a nonzero $\rho_1 \in \mathcal{I}$. Further, from (4) we obtain

$$\gamma_j \rho_j = \sum_{I_N, i_j = 0} (-q^2)^{j-1} \lambda_{I_N} \gamma_1 \gamma_2 \cdots \gamma_j \gamma_{j+1}^{i_{j+1}} \cdots \gamma_N^{i_N}.$$

If $\gamma_j \rho_j = 0$ then Proposition 2 gives $\lambda_{I_N} = 0$ for all I_N with $i_j = 0$. Hence $\rho_{j+1} := \rho_j$ fulfills our hypothesis ($*$) for $j + 1$. Otherwise $\gamma_j \rho_j \neq 0$ and the element $\rho_{j+1} := \gamma_j \rho_j$ satisfies ($*$) for $j + 1$. By induction it follows that ($*$) is valid for $j = n + 1$.

Now we prove by induction that

($**$) for all $j \in \{1, 2, \dots, n\}$ there exists $\rho_j = \sum_{I_N} \lambda_{I_N} \gamma^{I_N} \in \mathcal{I}$ such that $\lambda_{I_N} \neq 0$ implies $i_k = 1$ for $k \leq n$ and for $k > j'$.

By ($*$) this is true for $j = 1$. Further, we obtain

$$\rho_j \gamma_{j'} = \sum_{I_N, i_{j'} = 0} (-q^2)^{j-1} \lambda_{I_N} \gamma_1^{i_1} \cdots \gamma_{j'-1}^{i_{j'-1}} \gamma_{j'} \gamma_{j'+1} \cdots \gamma_N.$$

If $\rho_j \gamma_{j'} = 0$ then Proposition 2 implies $\lambda_{I_N} = 0$ whenever $i_{j'} = 0$. Hence $\rho_{j+1} := \rho_j$ satisfies ($**$). Otherwise $\rho_{j+1} := \rho_j \gamma_{j'} \neq 0$ and it fulfills ($**$). By induction we get ($**$) for $j = n + 1$. This means that $\gamma_1 \gamma_2 \cdots \gamma_N \in \mathcal{I}$.

Now we prove by induction that $\gamma_1 \gamma_2 \cdots \gamma_{j'} \in \mathcal{I}$ for $j \leq n + 1$. Indeed, this is true for $j = 1$. Further, from (4) and $j \leq n$ we get

$$\begin{aligned} \gamma_1 \gamma_2 \cdots \gamma_{j'} \gamma_j &= \gamma_1 \gamma_2 \cdots \gamma_{j'-1} \left(-\gamma_j \gamma_{j'} + [2] \hat{q} \sum_{k=1}^{j-1} q^{2k-2j+2} \gamma_k \gamma_{k'} + c^2 q^{N-2j+1} [2] \right) \\ &= c^2 q^{N-2j+1} [2] \gamma_1 \gamma_2 \cdots \gamma_{j'-1}. \end{aligned}$$

This means that $\gamma_1 \gamma_2 \cdots \gamma_n \in \mathcal{I}$. Finally, we prove by inverse induction on j that $\rho_j := \gamma_1 \gamma_2 \cdots \gamma_j \in \mathcal{I}$ for $j = n, n-1, \dots, 0$. This is true for $j = n$. Of course, then $\rho_j\gamma_{j'} \in \mathcal{I}$. On the other hand,

$$\begin{aligned}\gamma_{j'}\rho_j &= (-q^2)^{j-1}\gamma_1\gamma_2\cdots\gamma_{j-1}\gamma_{j'}\gamma_j\\ &= (-q^2)^{j-1}\gamma_1\cdots\gamma_{j-1}\left(-\gamma_j\gamma_{j'} + [2]\hat{q}\sum_{k=1}^{j-1} q^{2k-2j+2}\gamma_k\gamma_{k'} + c^2q^{N-2j+1}[2]\right)\\ &= -(-q^2)^{j-1}\rho_j\gamma_{j'} + (-1)^{j-1}c^2q^{N-1}[2]\gamma_1\gamma_2\cdots\gamma_{j-1}.\end{aligned}$$

Hence $\rho_{j-1} \in \mathcal{I}$. For $j = 0$ we get $\rho_0 = 1 \in \mathcal{I}$ and hence $\mathcal{I} = \mathrm{Cl}_{q^2}(N, c^2)$. ■

For $N = 2n + 1$ let $\mathcal{I}_+$ and $\mathcal{I}_-$ denote the two-sided ideal of $\mathrm{Cl}_{q^2}(N, c^2)$ generated by the element ρ_+ and ρ_-, respectively, where

$$\rho_\pm = \gamma_1\cdots\gamma_n(\pm c + \gamma_{n+1})\gamma_{n+2}\cdots\gamma_N. \tag{11}$$

Proposition 6. *The algebra* $\mathrm{Cl}_{q^2}(2n+1, c^2)$ *is semisimple and splits into the direct sum of the two simple ideals* $\mathcal{I}_+$ *and* $\mathcal{I}_-$*. Further,* $\dim\mathcal{I}_+ = \dim\mathcal{I}_-$*.*

Proof. Let $\mathcal{I} \neq \{0\}$ be a two-sided ideal of $\mathrm{Cl}_{q^2}(2n + 1, c^2)$. Similarly to the first part of the proof of Proposition 5 one can show that there exist complex numbers α_1, α_2, $|\alpha_1|+|\alpha_2| > 0$, such that $\rho := \gamma_1\gamma_2\cdots\gamma_n(\alpha_1+\alpha_2\gamma_{n+1})\gamma_{n+2}\cdots\gamma_N \in \mathcal{I}$. Multiplying ρ by $(-q^{-2})^n\gamma_{n+1}$ from the left we obtain that

$$\rho' := \gamma_1\cdots\gamma_n\gamma_{n+1}(\alpha_1 + \alpha_2\gamma_{n+1})\gamma_{n+2}\cdots\gamma_N = \gamma_1\cdots\gamma_n(c^2\alpha_2 + \alpha_1\gamma_{n+1})\gamma_{n+2}\cdots\gamma_N$$

belongs to $\mathcal{I}$. If ρ and ρ' are linearly independent then $\mathcal{I}$ contains both $\mathcal{I}_+$ and $\mathcal{I}_-$. Otherwise $\alpha_1^2 = c^2\alpha_2^2$ and $\mathcal{I}$ contains at least one of the ideals $\mathcal{I}_+$ and $\mathcal{I}_-$. Assume $\mathcal{I}_\eta \subset \mathcal{I}$ with $\eta = +$ or $\eta = -$. Since $\mathrm{Cl}_{q^2}(2n+1, c^2)$ is a finite dimensional algebra over $\mathbb{C}$, its radical is nilpotent. Suppose that $\mathcal{I}_\eta$ is contained in the radical of $\mathrm{Cl}_{q^2}(2n + 1, c^2)$. Then ρ_η has to be nilpotent. Let us determine ρ_η^2. First, for $j = 1, \ldots , n$ we compute

$$\begin{aligned}&\gamma_1\cdots\gamma_n(\eta c + \gamma_{n+1})\gamma_{n+2}\cdots\gamma_{j'}\cdot\gamma_j\gamma_{j+1}\cdots\gamma_n(\eta c + \gamma_{n+1}) =\\ &\quad= \gamma_1\cdots\gamma_n(\eta c + \gamma_{n+1})\gamma_{n+2}\cdots\gamma_{j'-1}\Big(-\gamma_j\gamma_j' + [2]\hat{q}\sum_{k=1}^{j-1} q^{2k-2j+2}\gamma_k\gamma_{k'}\\ &\qquad + c^2q^{N-2j+1}[2]\Big)\gamma_{j+1}\cdots\gamma_n(\eta c + \gamma_{n+1})\\ &\quad= c^2q^{N-2j+1}[2]\gamma_1\cdots\gamma_n(\eta c + \gamma_{n+1})\gamma_{n+2}\cdots\gamma_{j'-1}\cdot\gamma_{j+1}\cdots\gamma_n(\eta c + \gamma_{n+1}).\end{aligned}$$

Further,

$$\gamma_1 \cdots \gamma_n (\eta c + \gamma_{n+1})^2 = \gamma_1 \cdots \gamma_n \left(c^2 + 2\eta c \gamma_{n+1} + \hat{q} \sum_{k=1}^{n} q^{2k-2n} \gamma_k \gamma_{k'} + c^2 \right)$$
$$= 2\eta c \gamma_1 \cdots \gamma_n (\eta c + \gamma_{n+1}).$$

This yields

$$\rho_\eta \gamma_1 \gamma_2 \cdots \gamma_n (\eta c + \gamma_{n+1}) = 2\eta c^{2n+1} q^{n(n+1)} [2]^n \gamma_1 \gamma_2 \cdots \gamma_n (\eta c + \gamma_{n+1}) \qquad (12)$$

and hence $\rho_\eta^2 = 2\eta c^{2n+1} q^{n(n+1)} [2]^n \rho_\eta$. Therefore ρ_η is not nilpotent. The only possibility is that the radical is $\{0\}$ and hence $\mathrm{Cl}_{q^2}(N, c^2)$ is semisimple.

Since $\dim \mathrm{Cl}_{q^2}(2n+1, c^2) = 2^{2n+1} \neq k^2$ for all $k \in \mathbb{N}$, $\mathrm{Cl}_{q^2}(2n+1, c^2)$ is not simple and $\mathcal{I}_\eta \neq \mathrm{Cl}_{q^2}(2n+1, c^2)$ for $\eta = +$ or $\eta = -$. Then the complement ideal $\mathrm{Cl}_{q^2}(N, c^2) \ominus \mathcal{I}_\eta$ contains $\mathcal{I}_{-\eta}$ and we get $\mathrm{Cl}_{q^2}(2n+1, c^2) \ominus (\mathcal{I}_+ \oplus \mathcal{I}_-) = \{0\}$. Hence $\mathrm{Cl}_{q^2}(2n+1, c^2) = \mathcal{I}_+ \oplus \mathcal{I}_-$. Then the fact that each nontrivial simple ideal contains $\mathcal{I}_+$ or $\mathcal{I}_-$ in turn implies that both $\mathcal{I}_+$ and $\mathcal{I}_-$ are simple ideals. Moreover, since the mapping $\gamma_i \mapsto \gamma_i$, $i \neq n+1$, $\gamma_{n+1} \mapsto -\gamma_{n+1}$ defines an algebra isomorphism of $\mathrm{Cl}_{q^2}(2n+1, c^2)$ and maps $\mathcal{I}_+$ to $\mathcal{I}_-$, both ideals have the same dimension. ■

For $N = 2n+1$ let $\nu \in \{1, -1\}$ and for $N = 2n$ let $\nu = 1$. Let $\varphi_N^\nu \in \mathrm{Cl}_{q^2}(N, c^2)$ denote the element $\varphi_N^\nu = (\nu c + \gamma_{n+1}) \gamma_{n+2} \gamma_{n+3} \cdots \gamma_N$ for $N = 2n+1$ and $\varphi_N^1 = \gamma_{n+1} \gamma_{n+2} \cdots \gamma_N$ for $N = 2n$.

Proposition 7. *The left ideals* $\mathcal{I}_{\mathrm{L}}^\nu := \mathrm{Cl}_{q^2}(N, c^2) \varphi_N^\nu$ *of* $\mathrm{Cl}_{q^2}(N, c^2)$ *are minimal. For all* ν *the set* $\{\gamma_1^{i_1} \gamma_2^{i_2} \cdots \gamma_n^{i_n} \varphi_N^\nu \mid i_j \in \{0,1\}, j = 1, \ldots, n\}$ *forms a vector space basis of* $\mathcal{I}_{\mathrm{L}}^\nu$.

Proof. By definition the given elements belong to the left ideal $\mathcal{I}_{\mathrm{L}}^\nu$. On the other hand, the equations (4) imply that $\gamma_k \varphi_N^\nu = 0$ for $k > k'$ and $\gamma_{n+1} \varphi_{2n+1}^\nu = \nu c \varphi_{2n+1}^\nu$. Hence each expression $\gamma_1^{i_1} \gamma_2^{i_2} \cdots \gamma_N^{i_N} \varphi_N^\nu$, $i_l \in \{0,1\}$ for $l = 1, 2, \ldots, N$, is a scalar multiple of $\gamma_1^{i_1} \gamma_2^{i_2} \cdots \gamma_n^{i_n} \varphi_N^\nu$. The linear independence of the given set follows immediately from Proposition 2. Finally, the dimension of the ideal $\mathcal{I}_{\mathrm{L}}^\nu$ is 2^n. By Propositions 5 and 6 and from $\dim \mathrm{Cl}_{q^2}(N, c^2) = 2^N$ it follows that minimal left ideals have dimension 2^n. Hence $\mathcal{I}_{\mathrm{L}}^\nu$ is minimal. ■

Similarly there exist canonical minimal right ideals $\mathcal{I}_{\mathrm{R}}^\nu$ of $\mathrm{Cl}_{q^2}(N, c^2)$ generated by $\psi_{2n}^1 = \gamma_1 \gamma_2 \cdots \gamma_n$ and $\psi_{2n+1}^\nu = \gamma_1 \gamma_2 \cdots \gamma_n (\nu c + \gamma_{n+1})$, respectively.

3 Spin representations of $U_{q^2}(\mathfrak{so}_N)$

Let $N \in \mathbb{N}$, $N \geq 3$, $N = 2n + \epsilon$, where $n \in \mathbb{N}$ and $\epsilon \in \{0, 1\}$. If not otherwise stated we use the conventions in [16]. However we set $q_i = q^{d_i}$, $i = 1, 2, \ldots, n$, where $d_i = 2$ for $i < n$ and $d_n = 2 - \epsilon$. In [16] the algebra $U_{q^2}(\mathfrak{so}_N)$ is defined by the generators E_i, F_i, K_i and K_i^{-1}, $i = 1, 2, \ldots, n$, and relations

$$K_iK_j = K_jK_i, \qquad K_iK_i^{-1} = K_i^{-1}K_i = 1, \tag{13}$$

$$K_iE_jK_i^{-1} = q_i^{a_{ij}}E_j, \qquad K_iF_jK_i^{-1} = q_i^{-a_{ij}}F_j, \tag{14}$$

$$E_iF_j - F_jE_i = \delta_{ij}\frac{K_i - K_i^{-1}}{q_i - q_i^{-1}}, \tag{15}$$

$$\sum_{r=0}^{1-a_{ij}} (-1)^r q_i^{r(n-r)} \left[\!\!\left[\begin{matrix} 1 - a_{ij} \\ r \end{matrix} \right]\!\!\right]_{q_i} E_i^{1-a_{ij}-r}E_jE_i^r = 0, \quad i \neq j, \tag{16}$$

$$\sum_{r=0}^{1-a_{ij}} (-1)^r q_i^{r(n-r)} \left[\!\!\left[\begin{matrix} 1 - a_{ij} \\ r \end{matrix} \right]\!\!\right]_{q_i} F_i^{1-a_{ij}-r}F_jF_i^r = 0, \quad i \neq j, \tag{17}$$

$i, j = 1, 2, \ldots, n$, where $\left[\!\!\left[\begin{smallmatrix} m \\ p \end{smallmatrix} \right]\!\!\right]_{q^2} = \left[\!\!\left[\begin{smallmatrix} m \\ p \end{smallmatrix} \right]\!\!\right]$, and $\left[\!\!\left[\begin{smallmatrix} m \\ p \end{smallmatrix} \right]\!\!\right]_q$ is defined in the same way as $\left[\!\!\left[\begin{smallmatrix} m \\ p \end{smallmatrix} \right]\!\!\right]$ but everywhere q^2 has to be replaced by q.

By [16, Proposition 6.5] there exists a unique Hopf algebra structure on $U_{q^2}(\mathfrak{so}_N)$ with coproduct Δ, counit ε and antipode S such that

$$\Delta(K_i) = K_i \otimes K_i, \qquad \Delta(K_i^{-1}) = K_i^{-1} \otimes K_i^{-1}, \tag{18}$$

$$\Delta(E_i) = 1 \otimes E_i + E_i \otimes K_i, \qquad \Delta(F_i) = K_i^{-1} \otimes F_i + F_i \otimes 1, \tag{19}$$

$$\varepsilon(K_i) = \varepsilon(K_i^{-1}) = 1, \qquad \varepsilon(E_i) = \varepsilon(F_i) = 0, \tag{20}$$

$$S(K_i) = K_i^{-1}, \quad S(K_i^{-1}) = K_i, \quad S(E_i) = -E_iK_i^{-1}, \quad S(F_i) = -K_iF_i \tag{21}$$

for $i = 1, 2, \ldots, n$.

For $i = 1, 2, \ldots, n$ and $j = 1, 2, \ldots, N$ set

$$\lambda_{i,j} := \begin{cases} q^{-2\delta^j_i + 2\delta^j_{i+1} - 2\delta^j_{i'-1} + 2\delta^j_{i'}}, & i = 1, 2, \ldots, n-1, \\ q^{-2\delta^j_n + 2\delta^j_{n+2}}, & N = 2n+1, i = n, \\ q^{-2\delta^j_{n-1} - 2\delta^j_n + 2\delta^j_{n+1} + 2\delta^j_{n+2}}, & N = 2n, i = n. \end{cases} \tag{22}$$

Observe that $\lambda_{i,j}\lambda_{i,j'} = 1$ for all i, j and

$$q_i^{a_{ij}} = \begin{cases} \lambda_{i,j+1}\lambda_{i,j'}, & i \leq n, j < n, \\ \lambda_{i,n+1}\lambda_{i,n+2}, & i \leq n, j = n. \end{cases} \tag{23}$$

From [16, Section 8.4.1] we recall the formulas for the vector representation T_1 of $U_{q^2}(\mathfrak{so}_N)$ with highest weight $(1+\delta_{N3}, 0, \dots, 0)$ and highest weight vector $\mathbf{e}_N = \gamma_N$ with respect to the simple roots $\alpha_1, \alpha_2, \dots, \alpha_n$.

$$\begin{gathered}
T_1(E_i K_i^{-1}) = q^2 E_{i+1,i} - q^2 E_{i',i'-1}, \quad T_1(F_i) = E_{i,i+1} - E_{i'-1,i'}, \\
T_1(K_i) = D_i^{-1} D_{i+1} D_{i'-1}^{-1} D_{i'} \quad \text{for } N = 2n+\epsilon,\ i = 1, 2, \dots, n-1, \\
T_1(E_n K_n^{-1}) = q'(q^2 E_{n+1,n} - q E_{n+2,n+1}), \quad T_1(F_n) = q'(q^2 E_{n,n+1} - q^{-1} E_{n+1,n+2}), \\
T_1(K_n) = D_n^{-1} D_{n+2} \quad \text{for } N = 2n+1, \\
T_1(E_n K_n^{-1}) = q^2 E_{n+1,n-1} - q^2 E_{n+2,n}, \quad T_1(F_n) = E_{n-1,n+1} - E_{n,n+2}, \\
T_1(K_n) = D_{n-1}^{-1} D_n^{-1} D_{n+1} D_{n+2} \quad \text{for } N = 2n.
\end{gathered} \tag{24}$$

Here we used the notation $q' = [2]^{1/2}$ and $D_j = \sum_{k=1}^{N} q^{2\delta_j^k} E_{k,k}$, $j = 1, 2, \dots, n$, and $E_{k,l}$, $k, l = 1, 2, \dots, N$, are the matrix units. The aim of this section is to prove the following theorems.

Theorem 8. *There exists an algebra map $\pi : U_{q^2}(\mathfrak{so}_N) \to \mathrm{Cl}_{q^2}(N, c^2)$, $N = 2n+\epsilon$, such that*

$$\pi(K_i) = \sum_{l=0}^{i+1} \sum_{\substack{1 \le j_1 < j_2 < \\ \cdots < j_l \le i+1}} \left(\prod_{r=1}^{l} \frac{\lambda_{i,j_r} - q^{4l-4r}}{c^2 [2] q^{N+1-2j_r}} \right) \gamma_{j_1} \gamma_{j_2} \cdots \gamma_{j_l} \gamma_{j_l'} \cdots \gamma_{j_1'}, \tag{25}$$

$$\pi(K_i^{-1}) = \sum_{l=0}^{i+1} \sum_{\substack{1 \le j_1 < j_2 < \\ \cdots < j_l \le i+1}} \left(\prod_{r=1}^{l} \frac{\lambda_{i,j_r}^{-1} - q^{4l-4r}}{c^2 [2] q^{N+1-2j_r}} \right) \gamma_{j_1} \gamma_{j_2} \cdots \gamma_{j_l} \gamma_{j_l'} \cdots \gamma_{j_1'}, \tag{26}$$

$$\pi(E_i K_i^{-1}) = \frac{q^{2i+1-N}}{c^2 [2]} \sum_{l=0}^{i-1} \sum_{\substack{1 \le j_1 < j_2 < \\ \cdots < j_l \le i-1}} \left(\prod_{r=1}^{l} \frac{1 - q^{4r}}{c^2 [2] q^{N+1-2j_r}} \right) \gamma_{j_1} \cdots \gamma_{j_l} \gamma_{i+1} \gamma_{i'} \gamma_{j_l'} \cdots \gamma_{j_1'}, \tag{27}$$

$$\pi(F_i) = \frac{q^{2i+1-N}}{c^2 [2]} \sum_{l=0}^{i-1} \sum_{\substack{1 \le j_1 < j_2 < \\ \cdots < j_l \le i-1}} \left(\prod_{r=1}^{l} \frac{1 - q^{4r}}{c^2 [2] q^{N+1-2j_r}} \right) \gamma_{j_1} \cdots \gamma_{j_l} \gamma_i \gamma_{i'-1} \gamma_{j_l'} \cdots \gamma_{j_1'} \tag{28}$$

for $i = 1, 2, \dots, n-1$,

$$\pi(K_n) = q^{2-\epsilon} \sum_{l=0}^{n} \sum_{\substack{1 \le j_1 < j_2 < \\ \cdots < j_l \le n}} \left(\prod_{r=1}^{l} \frac{\lambda_{n,j_r} - q^{4l-4r}}{c^2 [2] q^{N+1-2j_r}} \right) \gamma_{j_1} \gamma_{j_2} \cdots \gamma_{j_l} \gamma_{j_l'} \cdots \gamma_{j_1'}, \tag{29}$$

$$\pi(K_n^{-1}) = q^{-2+\epsilon} \sum_{l=0}^{n} \sum_{\substack{1\le j_1<j_2< \\ \cdots<j_l\le n}} \left(\prod_{r=1}^{l} \frac{\lambda_{n,j_r}^{-1} - q^{4l-4r}}{c^2[2]q^{N+1-2j_r}} \right) \gamma_{j_1}\gamma_{j_2}\cdots\gamma_{j_l}\gamma_{j_l'}\cdots\gamma_{j_1'} \tag{30}$$

for $N = 2n + \epsilon$,

$$\pi(E_nK_n^{-1}) = \frac{q'}{c^2[2]} \sum_{l=0}^{n-1} \sum_{\substack{1\le j_1<j_2< \\ \cdots<j_l\le n-1}} \left(\prod_{r=1}^{l} \frac{1 - q^{4r}}{c^2[2]q^{N+1-2j_r}} \right) \gamma_{j_1}\cdots\gamma_{j_l}\gamma_{n+1}\gamma_{n+2}\gamma_{j_l'}\cdots\gamma_{j_1'}, \tag{31}$$

$$\pi(F_n) = \frac{q^{-1}q'}{c^2[2]} \sum_{l=0}^{n-1} \sum_{\substack{1\le j_1<j_2< \\ \cdots<j_l\le n-1}} \left(\prod_{r=1}^{l} \frac{1 - q^{4r}}{c^2[2]q^{N+1-2j_r}} \right) \gamma_{j_1}\cdots\gamma_{j_l}\gamma_{n}\gamma_{n+1}\gamma_{j_l'}\cdots\gamma_{j_1'} \tag{32}$$

for $N = 2n + 1$ *and*

$$\pi(E_nK_n^{-1}) = \frac{q^{-1}}{c^2[2]} \sum_{l=0}^{n-2} \sum_{\substack{1\le j_1<j_2< \\ \cdots<j_l\le n-2}} \left(\prod_{r=1}^{l} \frac{1 - q^{4r}}{c^2[2]q^{N+1-2j_r}} \right) \gamma_{j_1}\cdots\gamma_{j_l}\gamma_{n+1}\gamma_{n+2}\gamma_{j_l'}\cdots\gamma_{j_1'}, \tag{33}$$

$$\pi(F_n) = \frac{q^{-1}}{c^2[2]} \sum_{l=0}^{n-2} \sum_{\substack{1\le j_1<j_2< \\ \cdots<j_l\le n-2}} \left(\prod_{r=1}^{l} \frac{1 - q^{4r}}{c^2[2]q^{N+1-2j_r}} \right) \gamma_{j_1}\cdots\gamma_{j_l}\gamma_{n-1}\gamma_{n}\gamma_{j_l'}\cdots\gamma_{j_1'}, \tag{34}$$

for $N = 2n$.

Theorem 9. *(i) The vector space* $V = \mathbb{C}\{\gamma_i \,|\, i = 1, 2, \ldots, N\} \subset \mathrm{Cl}_{q^2}(N, c^2)$ *is invariant under the left adjoint action* $\pi_{\mathrm{ad}} : U_{q^2}(\mathfrak{so}_N) \to \mathrm{End}(\mathrm{Cl}_{q^2}(N, c^2))$ *of* $U_{q^2}(\mathfrak{so}_N)$*, where*

$$(\pi_{\mathrm{ad}} f)(v) = \pi(f_{(1)})\, v\, \pi(S(f_{(2)})), \qquad f \in U_{q^2}(\mathfrak{so}_N), \quad v \in \mathrm{Cl}_{q^2}(N, c^2). \tag{35}$$

The representation $\pi_{\mathrm{ad}} : U_{q^2}(\mathfrak{so}_N) \to \mathrm{End}(V)$ *of* $U_{q^2}(\mathfrak{so}_N)$ *is isomorphic to the vector representation with highest weight* $(1 + \delta_{N3}, 0, \ldots, 0)$ *and highest weight vector* γ_N*.*
(ii) For $N = 2n + 1$ *both mappings* $\pi^\nu : U_{q^2}(\mathfrak{so}_{2n+1}) \to \mathrm{End}(\mathcal{I}_{\mathrm{L}}^\nu)$, $\nu = -1, 1$, $(\pi^\nu f)(\psi) := \pi(f)\psi$ *for* $f \in U_{q^2}(\mathfrak{so}_{2n+1})$, $\psi \in \mathcal{I}_{\mathrm{L}}^\nu$*, are isomorphic to the spin representation of* $U_{q^2}(\mathfrak{so}_{2n+1})$ *with highest weight* $(0, 0, \ldots, 0, 1)$ *and highest weight*

vector φ^{ν}_{2n+1}.

(iii) For $N = 2n$ *the mapping* $\pi^{\nu} : U_{q^2}(\mathfrak{so}_{2n}) \to \mathrm{End}(\mathrm{Cl}^{\nu}_{q^2}(2n, c^2)\varphi^1_{2n})$, *where* $\nu = +, -$, $(\pi^{\nu} f)(\psi) := \pi(f)\psi$ *for* $f \in U_{q^2}(\mathfrak{so}_{2n})$, $\psi \in \mathrm{Cl}^{\nu}_{q^2}(2n, c^2)\varphi^1_{2n}$, *is isomorphic to the spin representation of* $U_{q^2}(\mathfrak{so}_{2n})$ *with highest weight* $(0, 0, \dots, 0, 0, 1)$ *and* $(0, 0, \dots, 0, 1, 0)$ *and highest weight vector* φ^1_{2n} *and* $\gamma_n\varphi^1_{2n}$ *for* $\nu = +$ *and* $\nu = -$, *respectively.*

Remark. In Proposition 3.1.3 and Theorem 3.1.2 in [6] the existence of a homomorphism from the algebra $U_q(\mathfrak{L}^{\pm})$ (which is closely related to $U_{q^2}(\mathfrak{so}_N)$, see [16, Sect. 8.5]) to $\mathrm{Cl}_{q^2}(N, c^2)$ was proved. Theorem 8 gives explicit formulas in terms of the generators of $U_{q^2}(\mathfrak{so}_N)$. ■

In order to prove these assertions we have to study the structure of $\mathrm{Cl}_{q^2}(N, c^2)$ in more detail.

Lemma 10. *Let* $\mu_i \in \mathbb{C}$, $i = 1, 2, \dots, N$, *and let* $\mu_j = 0$ *for some* $j \in \{1, 2, \dots, N\}$.*Then there exists no* $z \in \mathrm{Cl}_{q^2}(N, c^2)$, $z \neq 0$, *such that* $z\gamma_i = \mu_i\gamma_i z$ *for all* i.

Proof. Suppose that $z \in \mathrm{Cl}_{q^2}(N, c^2)$, $z \neq 0$, such that $z\gamma_i = \mu_i\gamma_i z$ for $i = 1, 2, \dots, N$. This yields in turn $z\mathrm{Cl}_{q^2}(N, c^2) \subset \mathrm{Cl}_{q^2}(N, c^2)z$. Further, $\mu_j = 0$ gives $z\gamma_j = 0$ and $\mathrm{Cl}_{q^2}(N, c^2)z\gamma_j = 0$. Hence $\mathcal{J}\gamma_j = 0$, where $\mathcal{J}$ denotes the two-sided ideal of $\mathrm{Cl}_{q^2}(N, c^2)$ generated by z. Since $z \in \mathcal{J}$, we have $\mathcal{J} \neq \{0\}$. Further, $\gamma_j \neq 0$ and $\mathcal{J}\gamma_j = 0$ imply $1 \notin \mathcal{J}$ and therefore $\mathcal{J} \neq \mathrm{Cl}_{q^2}(N, c^2)$. By Propositions 5 and 6 we conclude that $N = 2n + 1$ and $\mathcal{J} = \mathcal{I}_{\eta}$, $\eta \in \{+, -\}$. The relation $\rho_{\eta} \in \mathcal{J}$ and equation (12) give $\rho := \gamma_1\gamma_2 \cdots \gamma_n(\eta c + \gamma_{n+1}) \in \mathcal{J}$. Hence $\rho\gamma_j = 0$ implies $j \leq n$. Applying the algebra antiautomorphism τ to (12), from $\tau(\rho_{\eta}) = \rho_{\eta}$, $\tau(\mathcal{I}_{\eta}) = \mathcal{I}_{\eta}$ we get $\rho' := (\eta c + \gamma_{n+1})\gamma_{n+2}\gamma_{n+3} \cdots \gamma_N \in \mathcal{J}$. Now $\rho'\gamma_j = 0$ is fulfilled only for $j \geq n + 2$. This is a contradiction. ■

Lemma 11. *Let* $k \in \mathbb{N}$, $k \leq n$, *and let* $\mu, \mu' \in \mathbb{C}^{\times}$. *If* $z \in \mathrm{Cl}_{q^2}(N, c^2)$, $z \neq 0$, *satisfies* $z\gamma_k = \mu\gamma_k z$ *and* $z\gamma_{k'} = \mu'\gamma_{k'}z$, *then* z *is homogeneous with respect to* ∂_k *and we have* $\partial_k(z) = 0$.

Proof. There exist unique elements $z^{(r)} \in \mathrm{Cl}_{q^2}(N, c^2)$, $r = -1, 0, 1$, such that $\partial_k(z^{(r)}) = r$ and $z = z^{(-1)} + z^{(0)} + z^{(1)}$. The action of the algebra automorphism $\gamma_i \mapsto q^{\delta^i_k - \delta^i_{k'}}\gamma_i$, $i = 1, 2, \dots, N$, and its square on the equation $z\gamma_k - \mu\gamma_k z = 0$

give

$$(q^{-1}z^{(-1)}+z^{(0)}+qz^{(1)})\gamma_k=\mu\gamma_k(q^{-1}z^{(-1)}+z^{(0)}+qz^{(1)}), \tag{36}$$
$$(q^{-2}z^{(-1)}+z^{(0)}+q^2z^{(1)})\gamma_k=\mu\gamma_k(q^{-2}z^{(-1)}+z^{(0)}+q^2z^{(1)}). \tag{37}$$

Together with $z\gamma_k=\mu\gamma_k z$ we obtain $z^{(r)}\gamma_k=\mu\gamma_k z^{(r)}$ for $r=-1,0,1$. Similarly, $z^{(r)}\gamma_{k'}=\mu'\gamma_{k'}z^{(r)}$ for $r=-1,0,1$. We will show that $z^{(-1)}=z^{(1)}=0$.

By Proposition 2, $z^{(-1)}$ can uniquely be written as $z^{(-1)}=\sum_{I_N}\lambda_{I_N}\gamma^{I_N}$. Since $\partial_k(z^{(-1)})=-1$, for all multiindices I_N with $\lambda_{I_N}\neq 0$ we have $i_k=0$, $i_{k'}=1$. Let us fix a multiindex J_N in the set of those I_N for which $\lambda_{I_N}\neq 0$ and the sum $j_1+\cdots+j_{k-1}$ is minimal, say m. Consider the element

$$\gamma_1^{1-j_1}\gamma_2^{1-j_2}\cdots\gamma_{k-1}^{1-j_{k-1}}\gamma_k(z^{(-1)}\gamma_k-\mu\gamma_k z^{(-1)}).$$

Since $\gamma_k^2=0$, the second summand vanishes. Using the relations (4) the first summand becomes

$$\begin{aligned}
&\gamma_1^{1-j_1}\gamma_2^{1-j_2}\cdots\gamma_{k-1}^{1-j_{k-1}}\gamma_k z^{(-1)}\gamma_k=\\
&\qquad=\gamma_1^{1-j_1}\gamma_2^{1-j_2}\cdots\gamma_{k-1}^{1-j_{k-1}}\gamma_k\sum_{I_N}\lambda_{I_N}\gamma_1^{i_1}\gamma_2^{i_2}\cdots\gamma_N^{i_N}\gamma_k=\\
&\qquad=\gamma_1^{1-j_1}\gamma_2^{1-j_2}\cdots\gamma_{k-1}^{1-j_{k-1}}\gamma_k\sum_{I_N}\delta_0^{i_k}\delta_1^{i_{k'}}(-q^2)^{i_{k'+1}+\cdots+i_N}\lambda_{I_N}\gamma_1^{i_1}\gamma_2^{i_2}\cdots\gamma_{k'-1}^{i_{k'-1}}\times\\
&\qquad\quad\times\left(-\gamma_k\gamma_{k'}+\hat{q}[2]\sum_{l=1}^{k-1}q^{2l-2k+2}\gamma_l\gamma_{l'}+c^2q^{N-2k+1}[2]\right)\gamma_{k'+1}^{i_{k'+1}}\cdots\gamma_N^{i_N}.
\end{aligned}$$

Since $\gamma_k^2=0$, the first summand inside the large brackets vanishes for all I_N. Recall that if $i_k=0$ and $i_{k'}=1$ then $i_1+i_2+\ldots+i_{k-1}\geq m$. Hence for the number of factors γ_t, $1\leq t<k$, in the second expression inside of the brackets we obtain

$$\begin{aligned}
&(1-j_1)+(1-j_2)+\ldots+(1-j_{k-1})+i_1+i_2+\cdots+i_{k-1}+1\geq\\
&\qquad\geq k-1-j_1-j_2-\cdots-j_{k-1}+m+1=k-1-m+m+1=k.
\end{aligned}$$

This means that these summands are zero as well. Therefore we are left with

$$\begin{aligned}
\gamma_1^{1-j_1}\gamma_2^{1-j_2}\cdots\gamma_{k-1}^{1-j_{k-1}}\gamma_k z^{(-1)}\gamma_k&=c^2q^{N-2k+1}[2]\sum_{I_N}\delta_0^{i_k}\delta_1^{i_{k'}}(-q^2)^{n(I_N,J_N)}\lambda_{I_N}\times\\
&\times\gamma_1^{1-j_1+i_1}\gamma_2^{1-j_2+i_2}\cdots\gamma_{k-1}^{1-j_{k-1}+i_{k-1}}\gamma_k\gamma_{k+1}^{i_{k+1}}\cdots\gamma_{k'-1}^{i_{k'-1}}\gamma_{k'+1}^{i_{k'+1}}\cdots\gamma_N^{i_N}
\end{aligned}$$

with well-defined integer numbers $n(I_N, J_N)$. The appearing nonzero summands are linearly independent because of Proposition 2. Then $z^{(-1)}\gamma_k - \mu\gamma_k z^{(-1)} = 0$ implies that the above sum must be zero. In particular the summand for $I_N = J_N$ is zero if and only if $\lambda_{J_N} = 0$ which is a contradiction. Hence $z^{(-1)} = 0$.

Observe that $\partial_k(\tau(z^{(r)})) = -r$ and

$$\tau(z)\gamma_k = (\mu')^{-1}\gamma_k\tau(z). \tag{38}$$

Replacing μ by $(\mu')^{-1}$ and $z^{(-1)}$ by $\tau(z^{(1)})$ in the above proof, we obtain $\tau(z^{(1)}) = 0$ and hence $z^{(1)} = 0$. Therefore $z = z^{(0)}$. ■

Lemma 12. *Let* $\mu_i \in \mathbb{C}^\times$, $i = 1, 2, \ldots, N$, *and let* $z \neq 0$ *be an element in* $\mathrm{Cl}_{q^2}(N, c^2)$ *such that* $z\gamma_i = \mu_i\gamma_i z$ *for all* $i = 1, 2, \ldots, N$. *Then* $\mu_i\mu_{i'} = 1$ *and* $\mu_j = 1$ *for* $j = j'$.

Proof. From Lemma 11 we conclude that $\partial_k(z) = 0$ for all $k = 1, 2, \ldots, n$. Hence $\tau(z) = z$. The equations $\mu_k\gamma_k z - z\gamma_k = 0$ and (38) give

$$0 = \mu_k\gamma_k z - \mu_k\gamma_k\tau(z) = z\gamma_k - \mu_k\mu_{k'}\tau(z)\gamma_k = (1 - \mu_k\mu_{k'})z\gamma_k. \tag{39}$$

Since $z\gamma_k \neq 0$ by Lemma 10, we get $\mu_k\mu_{k'} = 1$. Finally, if $N = 2n + 1$ then the elements z and γ_{n+1} commute by Proposition 4. ■

Lemma 13. *Let* $\mu = (\mu_1, \mu_2, \ldots, \mu_k) \in (\mathbb{C}^\times)^k$, $1 \le k \le n$. *Then the following assertions are equivalent:*
(i) $z\gamma_i = \mu_i\gamma_i z$, $\tau(z)\gamma_i = \mu_i\gamma_i\tau(z)$ *for* $i = 1, 2, \ldots, k$,
(ii) $\partial_i(z) = 0$ *and*

$$z\gamma_1\gamma_2\cdots\gamma_i = \mu_1\mu_2\cdots\mu_i\gamma_1\gamma_2\cdots\gamma_i z, \quad \tau(z)\gamma_1\gamma_2\cdots\gamma_i = \mu_1\mu_2\cdots\mu_i\gamma_1\gamma_2\cdots\gamma_i\tau(z)$$

for $i = 1, 2, \ldots, k$,
(iii) $\partial_i(z) = 0$ *and* $\gamma_1\gamma_2\cdots\gamma_{i-1}(z\gamma_i - \mu_i\gamma_i z) = \gamma_1\gamma_2\cdots\gamma_{i-1}(\tau(z)\gamma_i - \mu_i\gamma_i\tau(z)) = 0$ *for* $i = 1, 2, \ldots, k$.

Proof. Applying the algebra antiautomorphism τ of $\mathrm{Cl}_{q^2}(N, c^2)$ to the second equation of (i) and using $\tau^2 = \mathrm{id}$ the conclusion (i)→(iii) follows immediately from Lemma 11. For the proof of (iii)→(ii) we take the sequence of equations

$$\mu_1\mu_2\cdots\mu_j\gamma_1\gamma_2\cdots\gamma_j z' = \mu_1\mu_2\cdots\mu_{j-1}(\gamma_1\gamma_2\cdots\gamma_{j-1}z'\gamma_j) =$$
$$= \mu_1\cdots\mu_{j-2}(\gamma_1\cdots\gamma_{j-2}z'\gamma_{j-1})\gamma_j = \cdots = \mu_1\gamma_1 z'\gamma_2\cdots\gamma_j = z'\gamma_1\gamma_2\cdots\gamma_j,$$

where the equations of (iii) for $i = j, j-1, \ldots, 2, 1$ and $z' = z, \tau(z)$ are used.

We have to prove (ii)→(i). Suppose that $z \in \mathrm{Cl}_{q^2}(N, c^2)$, $z \neq 0$, fulfills (ii). We prove by induction that $z\gamma_i = \mu_i\gamma_i z$ and $\tau(z)\gamma_i = \mu_i\gamma_i\tau(z)$ for $i = 1, 2, \ldots, k$. Our induction assumptions are the equations in (ii) for fixed $i := m < k$ and the equations $z\gamma_j = \mu_j\gamma_j z$, $z\gamma_{j'} = \mu_j^{-1}\gamma_{j'}z$ for all $j = 1, 2, \ldots, m-1$. Observe that the last requirements are empty at the beginning $m = 1$ of the induction. We have to prove $z\gamma_m = \mu_m\gamma_m z$ and $\tau(z)\gamma_m = \mu_m\gamma_m\tau(z)$. Applying τ to the last equation we then obtain $z\gamma_{m'} = \mu_m^{-1}\gamma_{m'}z$ which is needed for the next induction step.

Now we prove not only $z\gamma_m = \mu_m\gamma_m z$ but all equations

$$z\gamma_1\gamma_2\cdots\gamma_j\gamma_m = \mu_1\mu_2\cdots\mu_j\mu_m\gamma_1\gamma_2\cdots\gamma_j\gamma_m z, \tag{40}$$

$j = 0, 1, 2, \ldots, m-1$ by inverse induction on j. For $j = m-1$ this is the induction hypothesis and for $j = 0$ the equation we want to prove. For the induction step we use the formula $z\gamma_{j'} = \mu_j^{-1}\gamma_{j'}z$ which is known since $j < m$. By (4) we compute

$$\begin{aligned}
z\gamma_1\gamma_2\cdots\gamma_{j-1}\gamma_m &= z\left(\frac{-q^{2j+1-N}}{[2]c^2}\gamma_1\gamma_2\cdots\gamma_j\gamma_m\gamma_{j'} - (-1)^j\frac{q^{1-N}}{[2]c^2}\gamma_{j'}\gamma_1\gamma_2\cdots\gamma_j\gamma_m\right)\\
&= \mu_1\mu_2\cdots\mu_j\mu_m\mu_j^{-1}\left(\frac{-q^{2j+1-N}}{[2]c^2}\gamma_1\gamma_2\cdots\gamma_j\gamma_m\gamma_{j'} - (-1)^j\frac{q^{1-N}}{[2]c^2}\gamma_{j'}\gamma_1\gamma_2\cdots\gamma_j\gamma_m\right)z\\
&= \mu_1\mu_2\cdots\mu_{j-1}\mu_m\gamma_1\gamma_2\cdots\gamma_{j-1}\gamma_m z.
\end{aligned}$$

Changing the role of z and $\tau(z)$ the remaining assertion $\tau(z)\gamma_m = \mu_m\gamma_m\tau(z)$ can be shown in the same way. ■

Taking Lemma 10 and Lemma 12 into account we define complex vector spaces $\mathcal{Z}_N^\mu$ for $\mu = (\mu_1, \mu_2, \ldots, \mu_n) \in (\mathbb{C}^\times)^n$. Set $\mu_{i'} = \mu_i^{-1}$ for $i \in \{1, 2, \ldots, N\}$, $i > i'$, and $\mu_i = 1$ for $i = i'$. Then we define

$$\mathcal{Z}_N^\mu = \{z \in \mathrm{Cl}_{q^2}(N, c^2) \mid z\gamma_i = \mu_i\gamma_i z, \quad i = 1, 2, \ldots, N\}. \tag{41}$$

Similarly to the proof of Lemma 11 one can show that $\mathcal{Z}_N^\mu$ splits into the direct sum $\mathcal{Z}_N^{\mu,0} \oplus \mathcal{Z}_N^{\mu,1}$ of ∂_0-homogeneous components of even and odd degree, respectively. Since there are no nonzero elements z of $\mathrm{Cl}_{q^2}^-(2n, c^2)$ such that $\partial_k(z) = 0$ for all $k = 1, 2, \ldots, n$, we have $\mathcal{Z}_{2n}^{\mu,1} = \{0\}$.

Lemma 14. *Let* $\mu = (\mu_1, \mu_2, \ldots, \mu_n) \in (\mathbb{C}^\times)^n$. *Then* $z \in \mathcal{Z}_N^\mu$ *if and only if* $\partial_k(z) = 0$ *and* $\gamma_1\gamma_2\cdots\gamma_{k-1}(z\gamma_k - \mu_k\gamma_k z) = 0$ *for all* $k = 1, 2, \ldots, n$.

Proof. The only if part of the lemma follows from Lemma 11 and the definition of $\mathcal{Z}^{\mu}_{N}$. On the other hand, suppose that $\partial_k(z) = 0$ and $\gamma_1\gamma_2\cdots\gamma_{k-1}(z\gamma_k - \mu_k\gamma_k z) = 0$ for all $k = 1, 2, \ldots, n$. Then $\tau(z) = z$ and hence we can apply Lemma 13(iii)→(i) with $k = n$. This means $z\gamma_i = \mu_i\gamma_i z$ and $z\gamma_{i'} = \mu_i^{-1}\gamma_{i'}z$. The remaining assertion $z\gamma_{n+1} = \gamma_{n+1}z$ follows from $\{z, \gamma_{n+1}\} \subset \widehat{\mathrm{Cl}}_{q^2}(2n+1, c^2)$ and Proposition 4. ■

Proposition 15. *Let $\mu = (\mu_1, \mu_2, \ldots, \mu_n) \in (\mathbb{C}^\times)^n$. The vector spaces $\mathcal{Z}^{\mu,0}_{2n}, \mathcal{Z}^{\mu,0}_{2n+1}$ and $\mathcal{Z}^{\mu,1}_{2n+1}$ are one-dimensional and generated by the element*

$$z_{\mu,\hat\eta} = \sum_{k=0}^{n} \sum_{\substack{1\le i_1<i_2<\\ \cdots<i_k\le n}} \left(\prod_{r=1}^{k} \frac{\mu_{i_r} - (-q^2)^{\hat\eta} q^{4k-4r}}{c^2[2]q^{N+1-2i_r}} \right) \gamma_{i_1}\gamma_{i_2}\cdots\gamma_{i_k}\gamma_{n+1}^{\hat\eta}\gamma_{i'_k}\cdots\gamma_{i'_2}\gamma_{i'_1},$$

where $\hat\eta = 0$ for $\mathcal{Z}^{\mu,0}_{2n}$ and $\mathcal{Z}^{\mu,0}_{2n+1}$, and $\hat\eta = 1$ for $\mathcal{Z}^{\mu,1}_{2n+1}$.

Corollary 16. *The elements $z_0 \in \mathrm{Cl}_{q^2}(2n, c^2)$ and $z_1 \in \mathrm{Cl}^{-}_{q^2}(2n+1, c^2)$, given by*

$$z_0 = \sum_{k=0}^{n} (-1)^k \prod_{l=1}^{k} \frac{q^{2l-2} + q^{-2l+2}}{q[2]c^2} \sum_{\substack{J_n\\ \ell(J_n)=k}} q^{2L(J_n)-k(2n-k+1)} \gamma_1^{j_1}\gamma_2^{j_2}\cdots\gamma_n^{j_n}\gamma_{n+1}^{j_n}\cdots\gamma_N^{j_1}, \tag{42}$$

$$z_1 = \frac{1}{c} \sum_{k=0}^{n} \prod_{l=1}^{k} \frac{q^{2l-1} + q^{-2l+1}}{q[2]c^2} \sum_{\substack{J_n\\ \ell(J_n)=k}} q^{2L(J_n)-k(2n-k+1)} \gamma_1^{j_1}\gamma_2^{j_2}\cdots\gamma_n^{j_n}\gamma_{n+1}\gamma_{n'}^{j_n}\cdots\gamma_N^{j_1}, \tag{43}$$

satisfy the equations $z_0\gamma_i = -\gamma_i z_0$ and $z_1\gamma_i = \gamma_i z_1$, respectively, for all $i = 1, 2, \ldots, N$. Moreover, they are unique with this property up to a scalar factor.

Proof of the proposition. Let $z \in \mathcal{Z}^{\mu,\hat\eta}_{N}$, $z \neq 0$. We use Lemma 14 to determine the explicit form of z. The requirement $\partial_k(z) = 0$ for all $k = 1, 2, \ldots, n$ is equivalent to the fact that z takes the form

$$z = \sum_{I_n} \lambda_{I_n} \gamma_1^{i_1}\gamma_2^{i_2}\cdots\gamma_n^{i_n}\gamma_{n+1}^{\hat\eta}\gamma_{n'}^{i_n}\cdots\gamma_N^{i_1}.$$

If I_n is a multiindex with $i_k = 0$ for some $k \in \{1, 2, \ldots, n\}$, then let I_n^k denote the multiindex $(j_1, j_2, \ldots, j_n)$ such that $j_l = i_l + \delta_{lk}$, $l = 1, 2, \ldots, n$. Let us compute the left hand side of the remaining equations in Lemma 14. By (4) we obtain

$$\begin{aligned}
&\gamma_1\gamma_2\cdots\gamma_{k-1}(z\gamma_k-\mu_k\gamma_k z)= \qquad (44)\\
&\quad=\sum_{I_n,i_l=0\text{ for }l<k}\lambda_{I_n}\gamma_1\gamma_2\cdots\gamma_{k-1}(\gamma_k^{i_k}\gamma_{k+1}^{i_{k+1}}\cdots\gamma_n^{i_n}\gamma_{n+1}^{\hat\eta}\gamma_{n'}^{i_n}\cdots\gamma_{k'}^{i_k}\gamma_k\\
&\qquad\qquad-\mu_k\gamma_k\gamma_k^{i_k}\gamma_{k+1}^{i_{k+1}}\cdots\gamma_n^{i_n}\gamma_{n+1}^{\hat\eta}\gamma_{n'}^{i_n}\cdots\gamma_{k'}^{i_k})\\
&\quad=\sum_{I_n,i_l=0\text{ for }l\le k}\gamma_1\gamma_2\cdots\gamma_{k-1}(\lambda_{I_n^k}\gamma_k\gamma_{k+1}^{i_{k+1}}\cdots\gamma_{k'-1}^{i_{k+1}}\gamma_{k'}\gamma_k+\lambda_{I_n}\gamma_{k+1}^{i_{k+1}}\cdots\gamma_{k'-1}^{i_{k+1}}\gamma_k\\
&\qquad\qquad-\mu_k\lambda_{I_n}\gamma_k\gamma_{k+1}^{i_{k+1}}\cdots\gamma_{k'-1}^{i_{k+1}})\\
&\quad=\sum_{I_n,i_l=0\text{ for }l\le k}\gamma_1\gamma_2\cdots\gamma_{k-1}(\lambda_{I_n^k}c^2q^{N-2k+1}[2]+\lambda_{I_n}(-q^2)^{2i_{k+1}+2i_{k+2}+\cdots+2i_n+\hat\eta}\\
&\qquad\qquad-\mu_k\lambda_{I_n})\gamma_k\gamma_{k+1}^{i_{k+1}}\cdots\gamma_{k'-1}^{i_{k+1}}.
\end{aligned}$$

By Proposition 2 the summands with nonzero coefficient in the last expression are linearly independent. Hence $\gamma_1\gamma_2\cdots\gamma_{k-1}(z\gamma_k-\mu_k\gamma_k z)=0$ if and only if

$$\lambda_{I_n^k}=\frac{\mu_k-(-q^2)^{2i_{k+1}+2i_{k+2}+\cdots+2i_n+\hat\eta}}{c^2q^{N-2k+1}[2]}\lambda_{I_n} \qquad (45)$$

for all $k=1,2,\dots,n$. This yields $z\in\mathbb{C}z_{\mu,\hat\eta}$. ■

Recall the definition of z_0 and z_1 in Corollary 16.

Corollary 17. *We have* $z_0^2=1$ *in* $\mathrm{Cl}_{q^2}(2n,c^2)$ *and* $z_1^2=1$ *in* $\mathrm{Cl}_{q^2}(2n{+}1,c^2)$.

Proof. By definition z_ϵ is a ∂_0-homogeneous element in $\mathrm{Cl}_{q^2}(2n{+}\epsilon,c^2)$. Hence $z_\epsilon^2\in\mathrm{Cl}_{q^2}^+(2n{+}\epsilon,c^2)$ and by Corollary 16 it belongs to the center of $\mathrm{Cl}_{q^2}(2n{+}\epsilon,c^2)$. By Proposition 15 with $\mu:=(1,1,\dots,1)$, this implies $z_\epsilon^2=\lambda 1$ for some $\lambda\in\mathbb{C}$. Multiplying this equation by φ_N^1 from the right and taking into account the formula $z_\epsilon\varphi_N^1=\varphi_N^1$ we conclude that $\lambda=1$. ■

Corollary 18. *The element* z_1 *acts (by multiplication) with the factor* η *on the left ideal* $\mathcal{I}_{\mathrm{L}}^\eta$ *of* $\mathrm{Cl}_{q^2}(2n+1,c^2)$. *The element* z_0 *acts with the factor* $+1$ *and* -1 *on the subspaces* $\mathrm{Cl}_{q^2}^+(2n,c^2)\varphi_{2n}^1$ *and* $\mathrm{Cl}_{q^2}^-(2n,c^2)\varphi_{2n}^1$ *of the left ideal* $\mathcal{I}_{\mathrm{L}}^1$ *of* $\mathrm{Cl}_{q^2}(2n,c^2)$, *respectively.*

Now we turn to the proofs of Theorem 8 and Theorem 9. Therein the following lemma plays a crucial role.

Lemma 19. *For $f = K_i, K_i^{-1}, E_iK_i^{-1}, F_i$, $i = 1, 2, \dots, n$, let $\tilde{f}$ denote the element of $\mathrm{Cl}_{q^2}(N, c^2)$ given by the expression at the right of $\pi(f)$ in Theorem 8. Then the elements $\tilde{f}$ satisfy the following equations in $\mathrm{Cl}_{q^2}(N, c^2)$:*

$$\tilde{K}_i\gamma_j = \lambda_{i,j}\gamma_j\tilde{K}_i, \qquad \widetilde{K_i^{-1}}\gamma_j = \lambda_{i,j}^{-1}\gamma_j\widetilde{K_i^{-1}}, \tag{46}$$

$$\widetilde{E_kK_k^{-1}}\gamma_j = \begin{cases} \gamma_j\widetilde{E_kK_k^{-1}} & \textit{for } j \notin \{k, k+1, k'-1, k'\}, \\ q^{-2}\gamma_j\widetilde{E_kK_k^{-1}} & \textit{for } j = k+1, k', \\ q^2\gamma_j\widetilde{E_kK_k^{-1}} + q^2\gamma_{j+1} & \textit{for } j = k, \\ q^2\gamma_j\widetilde{E_kK_k^{-1}} - q^2\gamma_{j+1} & \textit{for } j = k'-1, \end{cases} \tag{47}$$

$$\tilde{F}_k\gamma_j = \begin{cases} \gamma_j\tilde{F}_k & \textit{for } j \notin \{k, k+1, k'-1, k'\}, \\ q^2\gamma_j\tilde{F}_k & \textit{for } j = k, k'-1, \\ q^{-2}\gamma_j\tilde{F}_k + \gamma_{j-1} & \textit{for } j = k+1, \\ q^{-2}\gamma_j\tilde{F}_k - \gamma_{j-1} & \textit{for } j = k', \end{cases} \tag{48}$$

$$\widetilde{E_nK_n^{-1}}\gamma_j = \begin{cases} \gamma_j\widetilde{E_nK_n^{-1}} & \textit{for } j \notin \{n-1, n, n+1, n+2\}, \\ \gamma_j\widetilde{E_nK_n^{-1}} & \textit{for } j = n-1,\ N = 2n+1, \\ q^2\gamma_j\widetilde{E_nK_n^{-1}} + q^2q'\gamma_{j+1} & \textit{for } j = n,\ N = 2n+1, \\ \gamma_j\widetilde{E_nK_n^{-1}} - qq'\gamma_{j+1} & \textit{for } j = n+1,\ N = 2n+1, \\ q^{-2}\gamma_j\widetilde{E_nK_n^{-1}} & \textit{for } j = n+2,\ N = 2n+1, \\ q^2\gamma_j\widetilde{E_nK_n^{-1}} + q^2\gamma_{j+2} & \textit{for } j = n-1,\ N = 2n, \\ q^2\gamma_j\widetilde{E_nK_n^{-1}} - q^2\gamma_{j+2} & \textit{for } j = n,\ N = 2n, \\ q^{-2}\gamma_j\widetilde{E_nK_n^{-1}} & \textit{for } j = n+1, n+2,\ N = 2n, \end{cases} \tag{49}$$

$$\tilde{F}_n\gamma_j = \begin{cases} \gamma_j\tilde{F}_n & \textit{for } j \notin \{n-1, n, n+1, n+2\}, \\ \gamma_j\tilde{F}_n & \textit{for } j = n-1,\ N = 2n+1, \\ q^2\gamma_j\tilde{F}_n & \textit{for } j = n,\ N = 2n+1, \\ \gamma_j\tilde{F}_n + q'\gamma_{j-1} & \textit{for } j = n+1,\ N = 2n+1, \\ q^{-2}\gamma_j\tilde{F}_n - q^{-1}q'\gamma_{j-1} & \textit{for } j = n+2,\ N = 2n+1, \\ q^2\gamma_j\tilde{F}_n & \textit{for } j = n-1, n,\ N = 2n, \\ q^{-2}\gamma_j\tilde{F}_n + \gamma_{j-2} & \textit{for } j = n+1,\ N = 2n, \\ q^{-2}\gamma_j\tilde{F}_n - \gamma_{j-2} & \textit{for } j = n+2,\ N = 2n, \end{cases} \tag{50}$$

for all $i = 1, 2, \dots, n$ and $k = 1, 2, \dots, n-1$, where $q' = [2]^{1/2}$.

Proof. Equation (46) follows in turn from Proposition 15. Observe that by definition $\widetilde{E_iK_i^{-1}} = q^{\delta_{\epsilon 1}\delta_{in}}\tau(\tilde{F}_i)$ for all $i = 1, 2, \dots, n$. Moreover, applying τ to the equations (47), (48), (49) and (50) we obtain the formulas (48), (47), (50) and (49), respectively, for j' instead of j. Therefore it suffices to show (47)–(50) for $j \leq j'$. Let us prove (47) and (48). The proof of the equations (49) and (50) can be treated similarly.

If $j \in \{k+2, k+3, \dots, n+\epsilon\}$ then $\tilde{F}_k\gamma_j = \gamma_j\tilde{F}_k$ and $\widetilde{E_kK_k^{-1}}\gamma_j = \gamma_j\widetilde{E_kK_k^{-1}}$. Indeed, in both $\tilde{F}_k$ and $\widetilde{E_kK_k^{-1}}$ there are no summands containing factors γ_l, $k+1 < l < k'-1$. Moreover, in each summand there are as many factors γ_l, $l \leq k+1$, as factors γ_l, $l \geq k'-1$. From this and the first line of (4) it follows that each summand of $\tilde{F}_k$ and of $\widetilde{E_kK_k^{-1}}$ commutes with γ_j.

If $j < k$ then we use Lemma 13(i)⇔(iii) with $(\mu_1, \mu_2, \dots, \mu_j) = (1, 1, \dots, 1)$. Obviously, $\partial_i(\widetilde{E_kK_k^{-1}}) = \partial_i(\tilde{F}_k) = 0$ for $i \leq j$. Moreover,

$$
\begin{aligned}
&\gamma_1\gamma_2\cdots\gamma_{j-1}(\tilde{F}_k\gamma_j - \gamma_j\tilde{F}_k) = \\
&= \gamma_1\gamma_2\cdots\gamma_{j-1}\sum_{I_n, i_l=0 \text{ for } l<j} \lambda_{I_n}(\gamma_j^{i_j}\gamma_{j+1}^{i_{j+1}}\cdots\gamma_{k-1}^{i_{k-1}}\gamma_k\gamma_{k'-1}\gamma_{k'+1}^{i_{k-1}}\cdots\gamma_{j'-1}^{i_{j+1}}\gamma_{j'}^{i_j}\gamma_j \\
&\qquad\qquad - \gamma_j\gamma_j^{i_j}\gamma_{j+1}^{i_{j+1}}\cdots\gamma_{k-1}^{i_{k-1}}\gamma_k\gamma_{k'-1}\gamma_{k'+1}^{i_{k-1}}\cdots\gamma_{j'-1}^{i_{j+1}}\gamma_{j'}^{i_j}) \\
&= \gamma_1\gamma_2\cdots\gamma_{j-1}\sum_{I_n, i_l=0 \text{ for } l\leq j} (\lambda_{I_n^j}\gamma_j\gamma_{j+1}^{i_{j+1}}\cdots\gamma_{k-1}^{i_{k-1}}\gamma_k\gamma_{k'-1}\gamma_{k'+1}^{i_{k-1}}\cdots\gamma_{j'-1}^{i_{j+1}}\gamma_{j'}\gamma_j \\
&\qquad\qquad + \lambda_{I_n}\gamma_{j+1}^{i_{j+1}}\cdots\gamma_{k-1}^{i_{k-1}}\gamma_k\gamma_{k'-1}\gamma_{k'+1}^{i_{k-1}}\cdots\gamma_{j'-1}^{i_{j+1}}\gamma_j \\
&\qquad\qquad - \lambda_{I_n}\gamma_j\gamma_{j+1}^{i_{j+1}}\cdots\gamma_{k-1}^{i_{k-1}}\gamma_k\gamma_{k'-1}\gamma_{k'+1}^{i_{k-1}}\cdots\gamma_{j'-1}^{i_{j+1}}) \\
&= \sum_{I_n, i_l=0 \text{ for } l\leq j} \gamma_1\gamma_2\cdots\gamma_{j-1}(\lambda_{I_n^j}c^2q^{N-2j+1}[2] + \lambda_{I_n}(-q^2)^{2i_{j+1}+\cdots+2i_{k-1}+2} \\
&\qquad\qquad - \lambda_{I_n})\gamma_j\gamma_{j+1}^{i_{j+1}}\cdots\gamma_{k-1}^{i_{k-1}}\gamma_k\gamma_{k'-1}\gamma_{k'+1}^{i_{k-1}}\cdots\gamma_{j'-1}^{i_{j+1}}.
\end{aligned}
$$

The last expression vanishes because of $\lambda_{I_n^j} = \dfrac{1 - q^{4\ell(I_n)+4}}{c^2q^{N+1-2j}[2]}$ for those coefficients λ_{I_n} of $\tilde{F}_k$ for which $i_l = 0$ for $l \leq j$ and $l \geq k$. Hence $\gamma_1\gamma_2\cdots\gamma_{j-1}(\tilde{F}_k\gamma_j - \gamma_j\tilde{F}_k) = 0$. Similarly one proves $\gamma_1\gamma_2\cdots\gamma_{j-1}(\widetilde{E_kK_k^{-1}}\gamma_j - \gamma_j\widetilde{E_kK_k^{-1}}) = 0$. Hence, by Lemma 13 we get (47) and (48) for $j < k$.

In a completely similar way one checks that

$$\begin{gathered}\gamma_1\gamma_2\cdots\gamma_{k-1}(\widetilde{E_kK_k^{-1}}\gamma_k - q^2\gamma_k\widetilde{E_kK_k^{-1}} - q^2\gamma_{k+1}) = 0,\\ \gamma_1\gamma_2\cdots\gamma_{k-1}(\tilde{F}_k\gamma_{k+1} - q^{-2}\gamma_{k+1}\tilde{F}_k - \gamma_k) = 0.\end{gathered} \tag{51}$$

Multiplying these equations from the left and from the right by $\gamma_{j'}$, $j = k-1, k-2, \dots, 1$, respectively, using (47) and (48) for $j > k'$, and taking the appropriate linear combination of the resulting expressions, one can successively cancel the leading factors γ_j, $j = k-1, k-2, \dots, 1$ in (51). This proves (47) for $j = k$ and (48) for $j = k+1$.

Finally, by definition we have $\partial_{k+1}(\widetilde{E_kK_k^{-1}}) = \partial_k(\tilde{F}_k) = 1$. Hence

$$\partial_{k+1}(\widetilde{E_kK_k^{-1}}\gamma_{k+1} - q^{-2}\gamma_{k+1}\widetilde{E_kK_k^{-1}}) = \partial_k(\tilde{F}_k\gamma_k - q^2\gamma_k\tilde{F}_k) = 2.$$

This and Proposition 2 imply that (47) and (48) are fulfilled for $j = k+1$ and $j = k$, respectively. ■

Proof of Theorem 8. We have to check that π maps all relations (13)–(17) of $U_{q^2}(\mathfrak{so}_N)$ to zero. Instead of this we prove that the corresponding equivalent relations, where the elements $K_i, K_i^{-1}, E_iK_i^{-1}$ and F_i are involved, are mapped to zero by π.

Since $\pi(K_i)$ and $\pi(K_j^{-1})$, $i, j = 1, 2, \dots, n$ are elements of $\widehat{\mathrm{Cl}}_{q^2}(N, c^2)$, by Proposition 4 they commute. Moreover, $z := \pi(K_i)\pi(K_i^{-1}) - 1$ and $z' := \pi(K_i^{-1})\pi(K_i) - 1$ are even elements of $\mathrm{Cl}_{q^2}(N, c^2)$ commuting with all γ_k, $k = 1, 2, \dots, N$ by (46). From Proposition 15 we get $z = \lambda z_{\mu,0} = \lambda 1$ and $z' = \lambda' z_{\mu,0} = \lambda' 1$, where $\mu = (1, 1, \dots, 1)$, $\lambda, \lambda' \in \mathbb{C}$. Hence $(z - \lambda)\varphi_N^1 = (z' - \lambda')\varphi_N^1 = 0$. On the other hand, $\pi(K_i)\varphi_N^1 = q^{(2-\epsilon)\delta_{in}}\varphi_N^1$ and $\pi(K_i^{-1})\varphi_N^1 = q^{(\epsilon-2)\delta_{in}}\varphi_N^1$ and therefore $\lambda\varphi_N^1 = \lambda'\varphi_N^1 = 0$. This means $\lambda = \lambda' = 0$ and hence equation (13) is compatible with π.

The compatibility of equations (14) (the first has to be multiplied by K_j^{-1}) with π follows from (46) and (23).

Since $\partial_i(\pi(E_i)) = \partial_i(\pi(E_iK_i^{-1})) = -1$, we have $\partial_i(\pi(E_i)^2) = -2$ for all $i = 1, 2, \dots, n$. By Proposition 2 this means that $\pi(E_i)^2 = 0$ for all $i = 1, 2, \dots, n$. Further, $\partial_{i+1}(\pi(E_i)) = 1$ for $i \le n-1$ and $\partial_i(\pi(E_{i+1})) = \partial_{i+2}(\pi(E_i)) = 0$ for $i \le n-2$. Hence $\partial_i(\pi(E_i)\pi(E_{i+1})\pi(E_i)) = -2$ and $\partial_{i+2}(\pi(E_{i+1})\pi(E_i)\pi(E_{i+1})) = 2$ for $i \le n-2$. Therefore $\pi(E_i)\pi(E_{i+1})\pi(E_i) = \pi(E_{i+1})\pi(E_i)\pi(E_{i+1}) = 0$ for

$i \leq n-2$. Similarly,

$$\pi(E_{n-2})\pi(E_n)\pi(E_{n-2}) = \pi(E_n)\pi(E_{n-2})\pi(E_n) = 0, \quad N = 2n,$$
$$\pi(E_{n-1})\pi(E_n)\pi(E_{n-1}) = 0, \quad N = 2n+1.$$

This proves that (16) is compatible with π for $a_{ij} < 0$. By similar arguments we obtain the same for (17).

Finally we have to prove compatibility of the relations (15) for all $i, j = 1, 2, \ldots, n$ and (16), (17) for $a_{ij} = 0$ with π. In order to do this we recall from equations (47)–(50) that there exist complex numbers $\hat{a}^l_{kj}$, $\hat{b}^l_{kj}$, $k = 1, 2, \ldots, n$, $j, l = 1, 2, \ldots, N$, such that

$$\pi(E_k K_k^{-1})\gamma_j = \lambda_{k,j}^{-1}\gamma_j \pi(E_k K_k^{-1}) + \hat{a}^l_{kj}\gamma_l, \qquad \pi(F_k)\gamma_j = \lambda_{k,j}^{-1}\gamma_j \pi(F_k) + \hat{b}^l_{kj}\gamma_l$$

for all $k = 1, 2, \ldots, n$, $j = 1, 2, \ldots, N$. Then

$$\begin{aligned}
&\left(q_i^{-a_{ik}} \pi(E_i K_i^{-1})\pi(F_k) - \pi(F_k)\pi(E_i K_i^{-1}) - \delta_{ik}\frac{1 - \pi(K_i^{-1})^2}{q_i - q_i^{-1}} \right)\gamma_j = \\
&= q_i^{-a_{ik}} \pi(E_i K_i^{-1})(\lambda_{k,j}^{-1}\gamma_j \pi(F_k) + \hat{b}^l_{kj}\gamma_l) - \pi(F_k)(\lambda_{i,j}^{-1}\gamma_j \pi(E_i K_i^{-1}) + \hat{a}^l_{ij}\gamma_l) \\
&\quad - \delta_{ik}\gamma_j \frac{1 - \lambda_{i,j}^{-2}\pi(K_i^{-1})^2}{q_i - q_i^{-1}} \\
&= \lambda_{i,j}^{-1}\lambda_{k,j}^{-1}\gamma_j \left(q_i^{-a_{ik}} \pi(E_i K_i^{-1})\pi(F_k) - \pi(F_k)\pi(E_i K_i^{-1}) - \delta_{ik}\frac{1 - \pi(K_i^{-1})^2}{q_i - q_i^{-1}} \right) \\
&\quad + (q_i^{-a_{ik}}\lambda_{i,l}^{-1}\hat{b}^l_{kj} - \lambda_{i,j}^{-1}\hat{b}^l_{kj})\gamma_l \pi(E_i K_i^{-1}) + (q_i^{-a_{ik}}\lambda_{k,j}^{-1}\hat{a}^l_{ij} - \lambda_{k,l}^{-1}\hat{a}^l_{ij})\gamma_l \pi(F_k) \\
&\quad + \left(q_i^{-a_{ik}}\hat{b}^l_{kj}\hat{a}^m_{il} - \hat{a}^l_{ij}\hat{b}^m_{kl} + \delta_{ik}\delta_{jm}\frac{\lambda_{i,j}^{-1}\lambda_{k,j}^{-1} - 1}{q_i - q_i^{-1}} \right)\gamma_m
\end{aligned}$$

for all $i, k = 1, 2, \ldots, n$ and $j = 1, 2, \ldots, N$. Observe that $T_1(E_i K_i^{-1})\gamma_j = \hat{a}^l_{ij}\gamma_l$, $T_1(F_k)\gamma_j = \hat{b}^l_{kj}\gamma_l$ and $T_1(K_k)\gamma_j = \lambda_{k,j}\gamma_j$ for all $i, k = 1, 2, \ldots, n$ and $j = 1, 2, \ldots, N$. Since

$$T_1(q_i^{-a_{ik}} K_i^{-1} F_k - F_k K_i^{-1})\gamma_j = 0, \quad T_1(q^{-d_i a_{ik}} E_i K_i^{-1} K_k^{-1} - K_k^{-1} E_i K_i^{-1})\gamma_j = 0,$$
$$T_1(q_i^{-a_{ik}} E_i K_i^{-1} F_k - F_k E_i K_i^{-1} + \delta_{ik}(K_i^{-2} - 1)/(q_i - q_i^{-1}))\gamma_j = 0,$$

the last two lines in the above expression vanish. Together with $\operatorname{im}\pi \subset \mathrm{Cl}^+_{q^2}(N, c^2)$ we obtain

$$z_{ik} := q_i^{-a_{ik}} \pi(E_i K_i^{-1})\pi(F_k) - \pi(F_k)\pi(E_i K_i^{-1}) - \delta_{ik}\frac{1 - \pi(K_i^{-2})}{q_i - q_i^{-1}} \in \mathcal{Z}_N^{\mu,0},$$

where $\mu = \mu_{(i,k)} = (\mu_1, \mu_2, \dots, \mu_n)$, $\mu_l = \lambda_{i,l}^{-1}\lambda_{k,l}^{-1}$. Similarly one can prove that

$$\begin{aligned} z'_{ik} &:= \pi(E_iK_i^{-1})\pi(E_kK_k^{-1}) - \pi(E_kK_k^{-1})\pi(E_iK_i^{-1}) \in \mathcal{Z}_N^{\mu,0}, \\ z''_{ik} &:= \pi(F_i)\pi(F_k) - \pi(F_k)\pi(F_i) \in \mathcal{Z}_N^{\mu,0} \end{aligned}$$

for $a_{ik} = 0$ and the same $\mu = \mu_{(i,k)}$. Since $\tau(z_{ik}) = z_{ki}$ and $\tau(z'_{ik}) = z''_{ki}$, it suffices to show that $z_{ik} = 0$ for $i \le k$ and $z'_{ik} = 0$ for $i, k = 1, 2, \dots, n$, $a_{ik} = 0$. By Proposition 15 we obtain $z_{ik}, z'_{ik} \in \mathbb{C}z_{\mu,0}$, $\mu = \mu_{(i,k)}$. Obviously, $z_{\mu,0}\varphi_N^1 = 1$. Hence we only have to prove that $z_{ik}\varphi_N^1 = 0$ for $i \le k$ and $z'_{ik}\varphi_N^1 = 0$ for $i, k = 1, 2, \dots, n$, $a_{ik} = 0$. Since $\pi(E_iK_i^{-1})\varphi_N^1 = 0$ for all $i = 1, 2, \dots, n$, we get $z'_{ik}\varphi_N^1 = 0$. If $i \le k < n$ then $\pi(K_i^{-1})\varphi_N^1 = \varphi_N^1$ and $\pi(F_k) = 0$. Hence $z_{ik}\varphi_N^1 = 0$. If $i < k = n$ then $\partial_i(\pi(E_iK_i^{-1})\pi(F_n)) = -1 + \delta^i_{n-1}\delta_{\epsilon 0}$. Moreover, for $i = n-1, k = n$ we get $\partial_n(\pi(E_{n-1}K_{n-1}^{-1})\pi(F_n)) = 2$. Hence for $i < k = n$ all summands of z_{ik} vanish after multiplication by φ_N^1. We are left with the case $i = k = n$, $N = 2n + \epsilon$. Then $\pi(E_nK_n^{-1})\varphi_N^1 = 0$ and $\pi(K_n^{-1})\varphi_N^1 = q^{\epsilon-2}\varphi_N^1$. For $N = 2n$ we compute

$$\begin{aligned} z_{nn}\varphi_{2n}^1 &= \left(q^{-4}\pi(E_nK_n^{-1})\frac{q^{-1}}{c^2[2]}\gamma_{n-1}\gamma_n - \frac{1-q^{-4}}{q^2-q^{-2}}\right)\varphi_{2n}^1 \\ &= \left(\frac{q^{-5}}{c^2[2]}\frac{q^{-1}}{c^2[2]}\gamma_{n+1}\gamma_{n+2}\gamma_{n-1}\gamma_n - q^{-2}\right)\varphi_{2n}^1 \\ &= \left(\frac{q^{-6}}{c^4[2]^2}\gamma_{n+1}c^2q^{N-2n+3}[2]\gamma_n - q^{-2}\right)\varphi_{2n}^1 \\ &= \left(\frac{q^{-3}}{c^2[2]}c^2q^{N-2n+1}[2] - q^{-2}\right)\varphi_{2n}^1 = 0. \end{aligned}$$

Similarly, for $N = 2n+1$ we obtain

$$\begin{aligned} z_{nn}\varphi_{2n+1}^1 &= \left(q^{-2}\pi(E_nK_n^{-1})\frac{q^{-1}q'}{c^2[2]}\gamma_n\gamma_{n+1} - \frac{1-q^{-2}}{q-q^{-1}}\right)\varphi_{2n+1}^1 \\ &= \left(\frac{q^{-3}}{cq'}\pi(E_nK_n^{-1})\gamma_n - q^{-1}\right)\varphi_{2n+1}^1 = \left(\frac{q^{-3}}{cq'}\frac{q'}{c^2[2]}\gamma_{n+1}\gamma_{n+2}\gamma_n - q^{-1}\right)\varphi_{2n+1}^1 \\ &= \left(\frac{q^{-3}}{c^3[2]}\gamma_{n+1}c^2q^{N-2n+1}[2] - q^{-1}\right)\varphi_{2n+1}^1 = \left(\frac{q^{-1}}{c}\gamma_{n+1} - q^{-1}\right)\varphi_{2n+1}^1 = 0. \end{aligned}$$

■

Proof of Theorem 9. First let us prove (i). Since π_{ad} is a representation of $U_{q^2}(\mathfrak{so}_N)$ on $\mathrm{Cl}_{q^2}(N, c^2)$, it is sufficient to check the invariance of V under the action of the generators $f = K_i, K_i^{-1}, E_iK_i^{-1}, F_i$ of $U_{q^2}(\mathfrak{so}_N)$. By the definition of $U_{q^2}(\mathfrak{so}_N)$, formula (35) for these f takes the form

$$(\pi_{\rm ad}K_i)(\gamma_j)=\pi(K_i)\gamma_j\pi(K_i^{-1}),\quad (\pi_{\rm ad}K_i^{-1})(\gamma_j)=\pi(K_i^{-1})\gamma_j\pi(K_i),$$
$$(\pi_{\rm ad}E_iK_i^{-1})(\gamma_j)=\pi(E_iK_i^{-1})\gamma_j-\pi(K_i^{-1})\gamma_j\pi(K_i)\pi(E_iK_i^{-1}),$$
$$(\pi_{\rm ad}F_i)(\gamma_j)=\pi(F_i)\gamma_j-\pi(K_i^{-1})\gamma_j\pi(K_i)\pi(F_i)$$

for $i=1,2,\dots,n$ and $j=1,2,\dots,N$. From this and Lemma 19 the invariance of V under the action $\pi_{\rm ad}$ of $U_{q^2}(\mathfrak{so}_N)$ immediately follows. Moreover,

$$(\pi_{\rm ad}K_i)(\gamma_N)=\lambda_{i,N}\gamma_N=q^{2\delta_{i1}}\gamma_N,$$
$$(\pi_{\rm ad}E_i)(\gamma_N)=(\pi_{\rm ad}E_iK_i^{-1})((\pi_{\rm ad}K_i)(\gamma_N))=q^{2\delta_{i1}}(\pi_{\rm ad}E_iK_i^{-1})(\gamma_N)=0,$$

and therefore the N-dimensional representation $\pi_{\rm ad}$ of $U_{q^2}(\mathfrak{so}_N)$ is isomorphic to the vector representation of $U_{q^2}(\mathfrak{so}_N)$ with higest weight $(1+\delta_{N3},0,0,\dots,0)$ and highest weight vector γ_N.

Now we turn to (ii). Since $\mathcal{I}_{\rm L}^\nu$, $\nu\in\{+,-\}$, is a left ideal of $\mathrm{Cl}_{q^2}(2n+1,c^2)$, the mapping π^ν gives a representation of $U_{q^2}(\mathfrak{so}_{2n+1})$. The dimension of this representation is $\dim\mathcal{I}_{\rm L}^\nu=2^n$ by Proposition 7. On the other hand, from $\pi(E_iK_i^{-1})\varphi_{2n+1}^\nu=0$ and $\pi(K_i)\varphi_{2n+1}^\nu=q^{\delta_{in}}$ for all $i=1,2,\dots,n$ we obtain that the 2^n-dimensional representation of $U_{q^2}(\mathfrak{so}_{2n+1})$ with highest weight $(0,0,\dots,0,1)$ and highest weight vector φ_{2n+1}^ν is a subrepresentation of π^ν. From this the assertion follows.

Finally we prove (iii). Since $\pi(f)\in\mathrm{Cl}_{q^2}^+(2n,c^2)$ for all $f=K_i,K_i^{-1},E_iK_i^{-1},F_i$, we have $\pi(U_{q^2}(\mathfrak{so}_{2n}))\subset\mathrm{Cl}_{q^2}^+(2n,c^2)$. Moreover, φ_{2n}^1 is homogeneous with respect to the $\mathbb{Z}_2$-grading ∂_0 of $\mathrm{Cl}_{q^2}(2n,c^2)$, and hence $\mathrm{Cl}_{q^2}^\nu(2n,c^2)\varphi_{2n}^1$, $\nu=+,-$, is an invariant subspace under left multiplication by $\mathrm{Cl}_{q^2}^+(2n,c^2)$, and therefore by $U_{q^2}(\mathfrak{so}_{2n})$ as well. By Proposition 7 the dimension of the vector space $\mathrm{Cl}_{q^2}^\nu(2n,c^2)\varphi_{2n}^1$, $\nu=+,-$, is 2^{n-1}. Again we obtain

$$(\pi^+K_i)(\varphi_{2n}^1)=q^{2\delta_{in}}\varphi_{2n}^1,\quad(\pi^+E_i)(\varphi_{2n}^1)=\pi(E_i)\varphi_{2n}^1=\pi(E_iK_i^{-1})q^{2\delta_{in}}\varphi_{2n}^1=0$$

for $i=1,2,\dots,n$. Hence π^+ contains as a subrepresentation the 2^{n-1}-dimensional highest weight representation with highest weight $(0,\dots,0,1)$ and highest weight vector φ^1_{2n}. Further,

$$\begin{aligned}(\pi^- K_i)(\gamma_n\varphi^1_{2n}) &= q^{2\delta_{in}}\left(1+\frac{\lambda_{i,n}-1}{c^2[2]q^{N+1-2n}}\gamma_n\gamma_{n+1}\right)\gamma_n\varphi^1_{2n}\\ &= q^{2\delta_{in}}\gamma_n\left(1+\frac{\lambda_{i,n}-1}{c^2[2]q^{N+1-2n}}c^2q^{N-2n+1}[2]\right)\varphi^1_{2n}\\ &= q^{2\delta_{in}}\lambda_{i,n}\gamma_n\varphi^1_{2n} = q^{2\delta^i_{n-1}}\gamma_n\varphi^1_{2n},\\ (\pi^- E_iK_i^{-1})(\gamma_n\varphi^1_{2n}) &= \pi(E_iK_i^{-1})\gamma_n\varphi^1_{2n} = \delta_{in}\frac{q^{-1}}{c^2[2]}\gamma_{n+1}\gamma_{n+2}\gamma_n\varphi^1_{2n} = 0\end{aligned}$$

for $i = 1, 2, \dots, n$. Hence π^- contains as a subrepresentation the 2^{n-1}-dimensional highest weight representation with highest weight $(0, 0, \dots, 0, 1, 0)$ and highest weight vector $\gamma_n\varphi^1_{2n}$. This proves the theorem. ■

References

[1] Ablamowicz, R. and Fauser, B.: On the decomposition of Clifford algebras of arbitrary bilinear form. Preprint math.QA/9911180.

[2] Bautista, R., Criscuolo, A., Durdević, M., Rosenbaum, M. and Vergara, J.D.: Quantum Clifford algebras from spinor representations. J. Math. Phys. **37**, 5747–5775 (1996)

[3] Birman, J.S. and Wenzl, H.: Braids, link polynomials and a new algebra. Trans. Am. Math. Soc. **313**(1), 249–273 (1989)

[4] Brzeziński, T., Papaloucas, L.C. and Rembieliński, J.: Quantum Clifford algebras. In *Clifford algebras and their applications in mathematical physics*, vol. 55 ser. *Fund. Theories Phys.*, pp. 3–8, Dordrecht (1993). Kluwer Acad. Publ.

[5] Carow-Watamura, U., Schlieker, M. and Watamura, S.: $SO_q(N)$ covariant differential calculus on quantum space and quantum deformation of Schrödinger equation. Z. Phys. C. **49**, 439–446 (1991)

[6] Ding, J. and Frenkel, I.B.: Spinor and oscillator representation of quantum groups. In *Lie theory and geometry*, vol. 123 ser. *Progr. Math.*, pp. 127–165, Boston (1994). Birkhäuser

[7] Durdević, M. and Oziewicz, Z.: Clifford algebras and spinors for arbitrary braids. Preprint q-alg/9412002.

[8] Faddeev, L.D., Reshetikhin, N.Yu. and Takhtajan, L.A.: Quantization of Lie groups and Lie algebras. Algebra and Analysis **1**, 178–206 (1987)

[9] Fiore, G.: On q-deformations of Clifford algebras. Preprint math.QA/0002194.

[10] Hannabuss, K.C.: Bilinear forms, Clifford algebras, q-commutation relations, and quantum groups. J. Algebra **228**, 227–256 (2000)

[11] Hayashi, T.: q-analogues of Clifford and Weyl algebras — spinor and oscillator representations of quantum enveloping algebras. Commun. Math. Phys. **127**, 129–144 (1990)

[12] Hayashi, T.: Quantum deformations of classical groups. Publ. Res. Inst. Math. Sci. **28**, 57–81 (1992)

[13] Heckenberger, I.: Hodge and Laplace–Beltrami operators for bicovariant differential calculi on quantum groups. Compositio Mathematica, in press. Preprint math.QA/9902130.

[14] Heckenberger, I.: Spin geometry on quantum groups via covariant differential calculi. Preprint NTZ 17/2000, math.QA/0006226.

[15] Heckenberger, I. and Schüler, A.: Symmetrizer and antisymmetrizer of the Birman–Wenzl–Murakami algebras. Lett. Math. Phys. **50**, 45–51 (1999)

[16] Klimyk, A. and Schmüdgen, K.: *Quantum groups and their representations*. Springer-Verlag, Heidelberg (1997)

[17] Takeuchi, M.: Matric bialgebras and quantum groups. Israel J. Math. **72**(1–2), 349–383 (1990)

[18] Welk, M.: Covariant first order differential calculus on quantum Euclidean spheres. Preprint math.QA/0008183.

[19] Woronowicz, S.L.: Differential calculus on compact matrix pseudogroups (quantum groups). Commun. Math. Phys. **122**(1), 125–170 (1989)